

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification 7 :</b> <b>C08K 5/103</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/26285</b> <b>(43) International Publication Date:</b> 11 May 2000 (11.05.00)
<b>(21) International Application Number:</b> PCT/US99/24840 <b>(22) International Filing Date:</b> 25 October 1999 (25.10.99)  <b>(30) Priority Data:</b> 60/106,094 29 October 1998 (29.10.98) US Not furnished 18 October 1999 (18.10.99) US  <b>(71) Applicant:</b> PENRECO [US/US]; Suite 400, 1 Shell Plaza, 910 Louisiana, Houston, TX 77002 (US).  <b>(72) Inventors:</b> HEALY, Lin, Lu; 1114 Augusta Dr. No.8, Houston, TX 77057 (US). CUNNINGHAM, Jack, C., Jr.; 5622 Knobby Knoll, Houston, TX 77092 (US). MORRISON, David, S.; 10600 Six Pines Dr. #1324, The Woodlands, TX 77380 (US). SONG, Wei; Apt. 242B, 7315 Brompton, Houston, TX 77025 (US). BUTUC, Gina; 14335 Ella Blvd. #1303, Houston, TX 77014 (US).  <b>(74) Agent:</b> BAI, J., Benjamin; Jenkins & Gilchrist, P.C., Suite 1800, 1100 Louisiana, Houston, TX 77002 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> GEL COMPOSITIONS  <b>(57) Abstract</b>  Gelled ester compositions are provided. The gelled esters are obtained by mixing a suitable ester with one or more triblock copolymers, star polymers, radial polymers, multi-block copolymers, or mixtures thereof. Optionally, one or more diblock copolymers can be used in forming the gel compositions. In addition to gelled esters, gelled alcohols, gelled ethers, and gelled naturally-occurring fats and oils also are obtained by using one or more diblock copolymers, triblock copolymers, star polymers, radial polymers, multi-block copolymers, or mixtures thereof as a gelling agent. Butyl rubbers and alkyl galactomannan also can be used as a gelling agent. The gel compositions may be used to suspend various solids and/or liquids.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## **GEL COMPOSITIONS**

### **CROSS-REFERENCE TO RELATED APPLICATIONS**

This patent application claims priority to a previously filed U.S. provisional application Serial No. 60/106,094, filed October 29, 1998, entitled "Gel Composition," in the name of Lin Lu, Jack Cunningham, Jr., and David S. Morrison. The disclosure of the aforementioned provisional patent application is incorporated by reference in its entirety herein.

### **FIELD OF THE INVENTION**

This invention relates to gel compositions which include esters, ethers, alcohols, naturally-occurring fats and oils, or mixtures thereof.

### **BACKGROUND OF THE INVENTION**

Numerous gel compositions are known in the art. Some have proven to be a convenient and efficient vehicle or carrier for the application of various active ingredients to the skin. Such active ingredients include antiperspirants, deodorants, perfumes, sunscreens, cosmetics, emollients, insect repellants, medicaments, and the like. Products incorporating a gel composition and those made entirely from a gel composition may be in the form of a soft gel or a stick. Rubbing a soft gel or solid stick containing an effective amount of an active ingredient dissolved or dispersed therein against the skin causes transfer of the gel composition to the skin surface in a layer form, leaving the active ingredient within the layer on the desired skin surface.

1           For topical administration of various cosmetic and health and beauty materials to the skin,  
2           a gel composition preferably should have one or more of the following desired properties:  
3           transparency, compatibility with the active ingredient, controlled release of the active ingredient,  
4           minimization of skin irritation, and the ability to suspend organic and inorganic materials, such as  
5           color pigments, glitters, water, air, metal oxides, sunscreen active particulates, and fragrances.  
6           In sunscreen applications, it is desirable if the gel itself could act as a sunscreen active ingredient.  
7           Moreover, it should moisturize the skin and exhibit water wash-off resistance, but should not have  
8           significant syneresis. For industrial applications, a gel base may offer one or more of the  
9           following properties: suspension, moisture barrier, rheology, solvency, controlled release, wetting,  
10          self-emulsifying, etc. For example, in the paint industry the properties that are important are  
11          suspension and controlled release.

12          A gel composition typically is made by mixing one or more compounds to be gelled with  
13          a gelling agent. Known gelling agents include metal soaps, surfactants, homopolymers, ionic  
14          homopolymers and copolymers, fumed silica, natural derivatives of gums, waxes, clay, and so on.  
15          Common gelling agents for cosmetic oils are fatty acid soaps of lithium, calcium, sodium,  
16          aluminum, zinc, and barium. A number of homopolymers and copolymers, such as atactic  
17          ethylene-propylene copolymers, have been used to gel hydrocarbons. For example, gelled mineral  
18          oils have been available; however, gel compositions including an ester, ether, alcohol, or vegetable  
19          oil are less common. Because gel compositions including an ester, alcohol, ether, or naturally-  
20          occurring fat or oil may provide a better alternative to hydrocarbon gels, there is a need for  
21          exploring methods of making such gel compositions.

### SUMMARY OF THE INVENTION

The invention meets the aforementioned need in one or more of the following aspects.

In one aspect, the invention relates to a gel composition that includes an ester compound and a polymer compound selected from the group consisting of triblock copolymers, star polymers, radial polymers, multi-block copolymers, and combinations thereof. The gel composition optionally may include a diblock copolymer. When the gel composition includes a diblock copolymer, the gel composition preferably is substantially free of mineral oils. In some embodiments, the gel composition may further include a suspended component. In other embodiments, the gel composition may further include an active ingredient. The active ingredient includes, but is not limited to, sun screens, antiperspirants, deodorants, perfumes, cosmetics, emollients, insect repellants, pesticides, herbicides, fungicides, plasticizers, insecticides, and medicaments.

In another aspect, the invention relates to a gel composition which includes at least two components. The first component is a compound selected from the group consisting of alcohols, ethers, naturally-occurring fats and oils, and combinations thereof. The second component is a polymer compound selected from the group consisting of diblock copolymers, triblock copolymers, star polymers, radial polymers, multi-block copolymers, and combinations thereof. In some embodiments, suitable alcohols include, but are not limited to, octyl dodecanol and isostearyl alcohol. In other embodiments, suitable ethers include, but are not limited to, dicarylyl ether. In still other embodiments, suitable naturally-occurring fats and oils include, but are not limited to, linseed oil, soybean oil, sunflower seed oil, corn oil, sesame oil, olive oil, castor oil, coconut oil, palm oil, and peanut oil.

1           In yet another aspect, the invention relates to a gel composition which includes at least  
2 two components. The first component is a compound selected from the group consisting of  
3 esters, alcohols, ethers, naturally-occurring fats and oils, and combinations thereof. The second  
4 component is a polymer compound selected from the group consisting of alkyl galactomannan,  
5 polybutadiene, and combinations thereof.

6           In still another aspect, the invention relates to a method of making a gel composition. The  
7 method includes (a) mixing an ester compound with a polymer compound selected from the group  
8 consisting of triblock copolymers, star polymers, radial polymers, multi-block copolymers, and  
9 combinations thereof; (b) heating the mixture; (c) agitating the mixture until the mixture becomes  
10 homogeneous; and (d) cooling the mixture.

11           In still yet another aspect, the invention relates to a method of making a gel composition.  
12 The method includes (a) mixing an alcohol, an ether, or a naturally-occurring fat or oil with a  
13 polymer compound selected from the group consisting of diblock copolymers, triblock  
14 copolymers, star copolymers, radial polymers, multi-block copolymers, and combinations thereof;  
15 (b) heating the mixture; (c) agitating the mixture until the mixture becomes homogeneous; and  
16 (d) cooling the mixture.

17           In another aspect, the invention relates to a method of making a gel composition. The  
18 method includes (a) mixing an ester, an alcohol, an ether, or a naturally-occurring fat or oil with  
19 alkyl galactomannan, or polybutadiene; (b) heating the mixture; (c) agitating the mixture until the  
20 mixture becomes homogeneous; and (d) cooling the mixture.

21           Properties and advantages of the embodiments of the invention become apparent with the  
22 following description.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

Fig. 1 is a schematic illustrating the chain structure of a diblock copolymer used in embodiments of the invention.

Fig. 2 is a schematic illustrating the chain structure of a triblock copolymer used in embodiments of the invention.

Fig. 3 is a schematic illustrating the chain structure of a radial polymer used in embodiments of the invention.

Fig. 4 is a schematic illustrating the chain structure of a star polymer used in embodiments of the invention.

Figs. 5A-5B are schematics illustrating the chain structure of multi-block copolymers used in embodiments of the invention.

### **DESCRIPTION OF EMBODIMENTS OF THE INVENTION**

Embodiments of the invention are, in part, based on the discovery that esters, alcohols, ethers, and naturally-occurring fats and oils can be gelled by using selected polymeric gelling agents. For example, diblock copolymers, triblock copolymers, radial polymers, star polymers, multi-block copolymers, butyl rubbers, alkyl galactomannan, and mixtures thereof can be used as gelling agents to obtain gelled esters, gelled ethers, gelled alcohols, and gelled naturally-occurring fats and oils. In some embodiments, the gel compositions and products made therefrom are transparent or substantially transparent. In other embodiments, the gel compositions and products made therefrom are semi-transparent, hazy, or opaque. These gel compositions and products

1 made therefrom each have numerous cosmetic and industrial applications.

2 A gel refers to a two-phase colloidal system comprising a liquid and a solid in the form  
3 of thickened liquid, semi-solid or solid. A gel also can refer to a composition that is either  
4 physically cross-linked by virtue of entangled polymer chains, or chemically cross-linked by  
5 virtue of covalent bonds such that it swells, but does not dissolve, in the presence of liquid.  
6 A gel typically is obtained by use of a gelling agent. The term "polymer" used herein includes  
7 both homopolymer and copolymer. A homopolymer is a polymer obtained by polymerizing  
8 one type of monomer, whereas a copolymer is a polymer obtained by polymerizing two or  
9 more types of monomers. "Block copolymer" refers to a copolymer in which like monomer  
10 units occur in relatively long, alternate sequences on a chain.

11 The term "opaque" refers to the optical state of a medium whose molecular aggregation  
12 is such that light cannot pass through. Therefore, light transmission through an opaque medium  
13 is substantially close to zero. On the other hand, the term "transparent" refers to the optical state  
14 of a medium through which light can pass through so that an object can be seen through it. As  
15 defined, the term "transparent" includes any optical state which is not opaque. A medium is  
16 considered transparent even if only a small fraction of light passes through it. Thus, a clear gel  
17 and a translucent gel are considered transparent.

18 In some embodiments of the invention, a block copolymer capable of forming a three-  
19 dimensional network through physical cross-linking is used as the gelling agent. Suitable block  
20 copolymers include at least one rigid block and one elastomeric block. The rigid blocks of a  
21 block copolymer form rigid domains through which physical cross-linking may occur. The  
22 physical cross-linking via these rigid domains yields a continuous three-dimensional network.



1 In the presence of heat, shear, or solvent, the rigid domains soften and permit flow. Upon  
2 cooling, removal of shear, or solvent evaporation, the rigid domains reform and harden,  
3 locking the elastomeric network in place. Preferably, suitable block copolymers include  
4 diblock copolymers, triblock copolymers, radial polymers, star polymers, multi-block  
5 copolymers, and mixtures thereof.

6 The amount of a gelling agent may range from about 0.2% to about 80% by weight,  
7 depending on the desired properties of the resulting gel. Preferably, a gelling agent is present  
8 in the gel from about 1% to about 40% by weight. More preferably, a gelling agent is present  
9 in the gel about 5% to about 20% by weight. In embodiments where both a diblock copolymer  
10 and a triblock copolymer are used, the triblock copolymer may range from about 0.1% to about  
11 10%, and the diblock copolymer from about 1% to about 40%.

12 Fig. 1 illustrates the typical chain structure of a diblock copolymer. The polymer chain  
13 of the diblock copolymer includes two blocks: a rigid block and an elastomeric block. The rigid  
14 block is represented by diamonds. The elastomeric block is represented by circles. The rigid  
15 block typically is composed of polystyrene, polyethylene, polyvinylchloride, phenolics, and the  
16 like; the elastomeric block may be composed of, ethylene/butadiene copolymers, polyisoprene,  
17 polybutadiene, ethylene/propylene copolymers, ethylene-propylene/diene copolymers, and the like.  
18 As such, suitable diblock copolymers include, but are not limited to, styrene-ethylene/propylene  
19 copolymers, styrene-ethylene/butadiene copolymers, styrene-isoprene copolymers, styrene-  
20 butadiene copolymers.

21 Fig. 2 illustrates the chain structure of a triblock copolymer. As illustrated in Fig. 2, each  
22 polymer chain includes two rigid blocks at either end and a middle block which is elastomeric.

1 This is a preferred triblock copolymer structure, although a triblock copolymer with two  
2 elastomeric end blocks and a rigid middle block also can be used. Suitable triblock copolymers  
3 include, but are not limited to, styrene-ethylene/propylene-styrene copolymers, styrene-  
4 ethylene/butadiene-styrene copolymers, styrene-isoprene-styrene copolymers, and styrene-  
5 butadiene-styrene copolymers. Multi-block copolymers are similar to diblock copolymers or  
6 triblock copolymers, except that the multiple block copolymers include additional elastomeric  
7 blocks and/or rigid blocks as illustrated in Figs. 5A-5B.

8 In addition to the linear chain structure, branched homopolymers or copolymers also may  
9 be used. Figs. 3-4 illustrate the chain structure of a radial polymer and a star polymer. It should  
10 be noted that one or more functional groups may be grafted onto the chain of any of the  
11 aforementioned polymers. In other words, any of the above polymers may be modified by  
12 grafting. Suitable functional groups for grafting depend on the desired properties. For example,  
13 one or more ester groups, silane groups, silicon-containing groups, maleic anhydride groups,  
14 acrylamide groups, and acid groups may be grafted. In addition to grafting, the above polymers  
15 may be hydrogenated to reduce unsaturation before they are used as gelling agents.

16 Numerous commercially available block copolymers may be used in embodiments of the  
17 invention. For example, various grades of copolymers sold under the trade name of Kraton® from  
18 Shell Chemical Company can be used as a gelling agent. In addition, copolymers sold under the  
19 trade name of Vector® available from Dexco and Septon® from Kuraray also may be used. U.S.  
20 Patent Nos. 5,221,534, No. 5,578,089, and No. 5,879,694 disclose block copolymers which may  
21 be used in embodiments of the invention, the disclosures of the three patents are incorporated by  
22 reference in their entirety herein. Table 1 lists some commercially available block copolymers

which may be used in embodiments of the invention. It is noted that additional suitable block copolymers may include, but are not limited to, polystyrene/polyester, polyether/polyamide, polyether/polyester, polyester/polyamide, polyether/polyurethane, polyester/polyurethane, poly(ethylene oxide)/poly(propylene oxide), nylon/rubber, and polysiloxane/polycarbonate.

TABLE 1

Copolymer	Block Type	Polystyrene Content(%)	Comment
Kraton® G 1702	SEP	28	hydrogenated diblock
Kraton® G 1701	SEP	37	hydrogenated diblock
Kraton® G 1780	SEP	7	star polymer
Kraton® G 1650	SEBS	30	hydrogenated triblock
Kraton® G 1652	SEBS	30	hydrogenated triblock
Kraton® D 1101	SEBS+SEP	31	triblock and diblock mixture (85:15)
Kraton® D 1102	SEBS + SEP	28	triblock + diblock (85:15)
Kraton® D 1133	SEBS +SEP	35	triblock + diblock (66:34)
Kraton® FG 1901	SEBS	30	triblock (hydrogenated and functionally grafted with 1.7% of maleic anhydride.
Septon® 1001	SEP	35	Hydrogenated diblock
Vector® 6030	SEP	30	Unsaturated diblock
Vector® 8550	SBS	29	Unsaturated triblock
Vector® 2518P	SBS	31	Unsaturated triblock
Solprene® 1430	SB	40	Unsaturated diblock


Note: SEP denotes to styrene/ethylene/propylene copolymers  
 SEBS denotes to styrene/ethylene/butylene/styrene copolymers  
 SB denotes to styrene/butadiene copolymers  
 SBS denotes to styrene-butadiene-styrene copolymers

1        It should be noted that block copolymers are not the only gelling agents that can be  
2        used in embodiments of the invention. Other types of polymers also may be used.  
3        Homopolymers which are capable of effecting strong molecular interaction between polymeric  
4        chains can be used as gelling agents. One such example is butyl rubber, which can thicken oil  
5        due to its compatibility with oil and high molecular weight. Specifically, a polybutadiene  
6        polymer sold under the trademark of Solprene® S200, which is available from GIRSA  
7        Industrias Negromex, S.A.de C.V. (INSA), can be used as a gelling agent. Other  
8        homopolymers capable of forming hydrogen bonding may include polyamide, polyester, and  
9        polyolefin. Still another example is alkyl galactomannan, which is capable of forming strong  
10       hydrogen bonding. Alkyl galactomannan is available from Hercules Incorporated under the  
11       trade name of N-HANCE® AG50 and AG200. Alkyl galactomannan is a polymer obtained by  
12       modifying natural products. It is especially effective in gelling esters and naturally-occurring  
13       fats and oils.


14       In accordance with embodiments of the invention, a gelled ester composition is  
15       obtained by gelling an ester compound with a triblock copolymer, a star polymer, a radial  
16       polymer, a multi-block copolymer, or mixtures thereof. Optionally, the gelled ester  
17       composition may further include one or more diblock copolymers. When a diblock  
18       copolymer is used along with one of triblock copolymers, star polymers, radial polymers, and  
19       multi-block copolymers, the resulting gel composition is substantially free of mineral oils.  
20       Suitable esters also may be gelled by alkyl galactomannan, polybutadiene, or other  
21       aforementioned polymers.

22       Any ester compound may be used in embodiments of the invention to obtain a gelled

ester composition. An ester is defined as a compound that includes one or more carboxylate groups:  $R-COO-$ , where R is hydrogen, hydrocarbyl, phenyl, methoxyphenyl, alkylphenyl, substituted alkyl, substituted phenyl, or other organic radicals. Suitable esters include monoesters, diesters, triesters, etc. For example, one class of suitable esters that can be gelled is represented by the following formulas:




and




where  $n=1, 2, 3$ , and  $4$ , and  $R_1$  includes hydrogen, hydrocarbyl, phenyl, methoxyphenyl, alkylphenyl, substituted alkyl, and substituted phenyl; and  $R_2$  includes hydrogen, hydrocarbyl, phenyl, methoxyphenyl, alkylphenyl, substituted alkyl, substituted phenyl, alkylene, phenylene, substituted alkylene, substituted phenylene, etc. It is noted that a suitable group for  $R_2$  depends on whether  $n$  is  $1, 2, 3$ , or  $4$ .

Another class of suitable esters that may be gelled in embodiments of the invention is represented by the following formula:



1 where  $R_3$  includes alkylene, phenylene, substituted alkylene, and substituted phenylene.

2 Still another class of suitable esters that may be gelled in embodiments of the invention  
3 is represented by the following formula:









11 where  $R_4$ ,  $R_5$ , and  $R_6$  individually include alkylene, phenylene, substituted alkylene, and  
12 substituted phenylene;  $R_7$ ,  $R_8$ , and  $R_9$  individually include hydrogen, hydrocarbyl, phenyl,  
13 methoxyphenyl, alkylphenyl, substituted alkyl, and substituted phenyl.

14 Preferred esters and their chemical formulas are listed in Table 2.

15 TABLE 2

16 Chemical Name	Chemical Formula
17 Isononyl Isononanoate	$  \begin{array}{c}  \text{O} \\  \parallel \\  C_8H_{17} - C - O - C_9H_{19}  \end{array}  $
18 Isopropyl Palmitate	$  \begin{array}{c}  \text{O} \\  \parallel \\  C_{15}H_{31} - C - O - C_3H_7  \end{array}  $

Chemical Name	Chemical Formula
1 C <sub>12-15</sub> alkyl benzoate	
2 Myristyl Neopentanoate	
3 Tridecyl Salicylate	
4 Octyl Methoxycinnamate	
5 Propylene glycol 6 dicaprylate/caprate	
7 Pentaerythrityl tetraisostearate	 <p data-bbox="771 1722 1323 1753">(R represents fatty acid radicals derived from coconut oil.)</p>

Chemical Name	Chemical Formula
Trimethylolpropane triisostearate	$  \begin{array}{c}  \text{CH}_2-\text{O}-\overset{\text{O}}{\parallel}{\text{C}}-\text{R} \\    \\  \text{CH}_3-\text{CH}_2-\text{C}-\text{CH}_2-\text{O}-\overset{\text{O}}{\parallel}{\text{C}}-\text{R} \\    \\  \text{CH}_2-\text{O}-\overset{\text{O}}{\parallel}{\text{C}}-\text{R}  \end{array}  $ <p>(R represents fatty acid radicals derived from coconut oil.)</p>
Glyceryl isostearate	$  \begin{array}{c}  \text{O} \\  \parallel \\  \text{C}_{17}\text{H}_{35}-\text{C}-\text{O}-\text{CH}_2-\text{CH}-\text{CH}_2\text{OH} \\    \\  \text{OH}  \end{array}  $
Diisononyl adipate	$  \begin{array}{c}  \text{O} \qquad \qquad \text{O} \\  \parallel \qquad \qquad \parallel \\  \text{C}_9\text{H}_{19}-\text{O}-\text{C}-(\text{CH}_2)_4-\text{C}-\text{O}-\text{C}_9\text{H}_{19}  \end{array}  $
Trioctyldodecyl citrate	$  \begin{array}{c}  \text{CH}_2-\text{O}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{R} \\    \\  \text{HO}-\text{C}-\text{C}-\text{O}-\text{R} \\    \qquad \qquad \parallel \\  \text{CH}_2-\text{O}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{R}  \end{array}  $ <p>(R is octyl dodecyl)</p>

Other suitable esters include, but are not limited to, the following compounds:

Acetylline Methylsilanol Mannuronate; Acetaminosalol; Acetylated Cetyl Hydroxyproline;

Acetylated Glycol Stearate; Acetylated Sucrose Distearate; Acetylmethionyl Methylsilanol



1 Elastinate; Acetyl Tributyl Citrate; Acetyl Triethyl Citrate; Acetyl Trihexyl Citrate; Aleurites  
2 Moluccana Ethyl Ester; Allethrins; Allyl Caproate; Amyl Acetate; Amyl Benzoate; Amyl  
3 Salicylate; Arachidyl Behenate; Arachldyl Glycol Isostearate; Arachldyl Propionate; Ascorbyl  
4 Dipalmitate; Ascorbyl Palmitate; Ascorbyl Stearate; Aspartame; Batyl Isostearate; Batyl  
5 Stearate; Bean Palmitate; Behenyl Beeswax; Behenyl Behenate; Behenyl Benzoate; Behenyl  
6 Erucate; Behenyl Isostearate; Behenyl/Isostearyl Beeswax; 1,2,4-Benzenetriacetate; Benzoin  
7 (Styrax Bensoin) Gum; Benzoxiquine; Benzyl Acetate; Benzyl Benzoate; Benzyl Cinnamate;  
8 Benzyl Hyaluronate; Benzyl Laurate; Benzyl Nicotinate; Benzylparaben; Benzyl Salicylate;  
9 Beta-Sitosteryl Acetate; Borago Officinalis Ethyl Ester; Butoxyethyl Acetate; Butoxyethyl  
10 Nicotinate; Butoxyethyl Stearate; Butyl Acetate; Butyl Acetyl Ricinoleate; Butyl Benzoate;  
11 Butyl Benzyl Phthalate; 2-t-Butylcyclohexyl Acetate; Butylene Glycol Dicaprylate/Dicaprate;  
12 Butylene Glycol Montanate; Butyl Ester of Ethylene/MA Copolymer; Butyl Ester of PVM/MA  
13 Copolymer; Butylglucoside Caprate; Butyl Isostearate; Butyl Lactate; Butyl Methacrylate;  
14 Butyl Myristate; Butyloctyl Beeswax; Butyloctyl Benzoate; Butyloctyl Candelillate; Butyloctyl  
15 Oleate; Butyloctyl Salicylate; Butyl Oleate; Butyl PABA; Butylparaben; Butyl Phthalyl Butyl  
16 Glycolate; Butyl Stearate; Butyl Thioglycolate; Butyroyl Trihexyl Citrate; C18-36 Acid Glycol  
17 Ester; C12-20 Acid PEG-8 Ester; Caffeine Benzoate; Calcium Pantetheine Sulfonate; Calcium  
18 Stearoyl Lactylate; C-18-28 Alkyl Acetate; C18-38 Alkyl Beeswax; C30-50 Alkyl Beeswax;  
19 C20-40 Alkyl Behenate; C18-38 Alkyl C24-54 Acid Ester; C-8 Alkyl Ethyl Phosphate; C18-38  
20 Alkyl Hydroxystearoyl Stearate; C12-13 Alkyl Lactate; C12-15 Alkyl Lactate; C12-13 Alkyl  
21 Octanoate; C12-15 Alkyl Octanoate; C12-15 Alkyl Salicylate; C18-36 Alkyl Stearate; C20-40  
22 Alkyl Stearate; C30-50 Alkyl Stearate; C40-60 Alkyl Stearate; Caproyl Ethyl Glucoside;

1 Capryloyl Salicylic Acid; Caprylyl Butyrate; Castor Oil Benzoate; C10-30  
2 Cholesterol/Lanosterol Esters; Cellulose Acetate; Cellulose Acetate Butyrate; Cellulose Acetate  
3 Propionate; Cellulose Acetate Propionate Carboxylate; Cetareth-7 Stearate; Cetearyl  
4 Behenate; Cetearyl Candelillate; Cetearyl Isononanoate; Cetearyl Octanoate; Cetearyl  
5 Palmitate; Cetearyl Stearate; Cetyl Acetate; Cetyl Acetyl Ricinoleate; Cetyl Caprylate; Cetyl  
6 C12-15-Pareth-9 Carboxylate; Cetyl Glycol Isostearate; Cetyl Isononanoate; Cetyl Lactate;  
7 Cetyl Laurate; Cetyl Myristate; Cetyl Octanoate; Cetyl Oleate; Cetyl Palmitate; Cetyl PCA;  
8 Cetyl PPG-2 Isodeceth-7 Carboxylate; Cetyl Ricinoleate; Cetyl Ricinoleate Benzoate; Cetyl  
9 Stearate; C16-20 Glycol Isostearate; C20-30 Glycol Isostearate; C14-16 Glycol Palmitate;  
10 Chimyl Isostearate; Chimyl Stearate; Chlorogenic Acids; Cholesteryl Acetate;  
11 Cholesteryl/Behenyl/Octyldecyl Lauroyl Glutamate; Cholesteryl Butyrate; Cholesteryl  
12 Dichlorobenzoate; Cholesteryl Hydroxystearate; Cholesteryl Isostearate; Cholesteryl Isostearyl  
13 Carbonate; Cholesteryl Lanolate; Cholesteryl Macadamiate; Cholesteryl Nonanoate;  
14 Cholesteryl/Octyldecyl Lauroyl Glutamate; Cholesteryl Oleate; Cholesteryl Stearate;  
15 Cinnamyl Acetate; Cinoxate; Citronellyl Acetate; Coco-Caprylate/Caprate; Coco Rapeseedate;  
16 Cocoyl Ethyl Glucoside; Copper PCA Methylsilanol; Corylus Avellanna Ethyl Ester; C12-15  
17 Pareth-9 Hydrogenated Tallowate; C11-15 Pareth-3 Oleate; C12-15 Pareth-12 Oleate; C11-15  
18 Pareth-3 Stearate; C11-15 Pareth-12 Stearate; Decyl Isostearate; Decyl Myristate; Decyl  
19 Oleate; Decyl Succinate; DEDM Hydantoin Dilaurate; Dextrin Behenate; Dextrin Laurate;  
20 Dextrin Myristate; Dextrin Palmitate; Dextrin Stearate; Diacetin; Dibutyl Adipate; Dibutyl  
21 Oxalate; Dibutyl Phthalate; Dibutyl Sebacate; Di-C12-15 Alkyl Adipate; Di-C12-15 Alkyl  
22 Fumarate; Di-C12-13 Alkyl Malate; Di-C12-13 Alkyl Tartrate; Di-C14-15 Alkyl Tartrate;

- 1 Dicapryl Adipate; Dicapryl Maleate; Dicetearyl Dimer Dilinoleate; Dicetyl Adipate; Dicetyl
- 2 Thiodipropionate; Dicoctyl Pentaerythrityl Distearyl Citrate; Diethoxyethyl Succinate; Diethyl
- 3 Acetyl Aspartate; Diethylaminoethyl Cocoate; Diethylaminoethyl PEG-5 Cocoate;
- 4 Diethylaminoethyl PEG-5 Laurate; Diethylaminoethyl Stearate; Diethyl Aspartate; Diethylene
- 5 Glycol Dibenzoate; Diethylene Glycol Diisononanoate; Diethylene Glycol Dioctanoate;
- 6 Diethylene Glycol Dioctanoate/Diisononanoate; Diethyl Glutamate; Diethyl Oxalate; Diethyl
- 7 Palmitoyl Aspartate; Diethyl Phthalate; Diethyl Sebacate; Diethyl Succinate; Digalloyl
- 8 Trioleate; Diglyceryl Stearate Malate; Dihexyl Adipate; Dihexyldecyl Lauroyl Glutamate;
- 9 Dihydroabietyl Behenate; Dihydroabietyl Methacrylate; Dihydrocholesteryl Butyrate;
- 10 Dihydrocholesteryl Isostearate; Dihydrocholesteryl Macadamiate; Dihydrocholesteryl
- 11 Nonanoate; Dihydrocholesteryl Octyldecanoate; Dihydrocholesteryl Oleate; Dihydrogenated
- 12 Palmoyl Hydroxyethylmonium Methosulfate; Dihydrogenated Tallow Phthalate;
- 13 Dihydrophytosteryl Octyldecanoate; Dihydroxyethylamino Hydroxypropyl Oleate;
- 14 Dihydroxyethyl Soyamine Dioleate; Diisobutyl Adipate; Diisobutyl Oxalate; Diisocetyl
- 15 Adipate; Diisodecyl Adipate; Diisopropyl Adipate; Diisopropyl Dimer Dilinoleate;
- 16 Diisopropyl Methyl Cinnamate; Diisopropyl Oxalate; Diisopropyl Sebacate;
- 17 Diisostearamidopropyl Epoxypropylmonium Chloride; Diisostearyl Trimethylolpropane
- 18 Siloxy Silicate; Diisostearyl Adipate; Diisostearyl Dimer Dilinoleate; Diisostearyl Fumarate;
- 19 Diisostearyl Glutarate; Diisostearyl Malte; Dilaureth-7 Citrate; Dilauryl Thiodipropionate;
- 20 Dimethicone Copolyol Acetate; Dimethicone Copolyol Adipate; Dimethicone Copolyol
- 21 Almondate; Dimethicone Copolyol Beeswax; Dimethicone Copolyol Behenate; Dimethicone
- 22 Copolyol Benzoate; Dimethicone Copolyol Borageate; Dimethicone Copolyol Cocoa Butterate;

1 Dimethicone Copolyol Dhupa Butterate; Dimethicone Copolyol Hydroxystearate; Dimethicone  
2 Copolyol Isostearate; Dimethicone Copolyol Kokum Butterate; Dimethicone Copolyol Lactate;  
3 Dimethicone Copolyol Laurate; Dimethicone Copolyol Mango Butterate; Dimethicone  
4 Copolyol Meadowfoamate; Dimethicone Copolyol Mohwa Butterate; Dimethicone Copolyol  
5 Octyldodecyl Citrate; Dimethicone Copolyol Olivatate; Dimethicone Copolyol Phthalate;  
6 Dimethicone Copolyol Sal Butterate; Dimethicone Copolyol Shea Butterate; Dimethicone  
7 Copolyol Stearate; Dimethicone Copoly Undecylenate; Dimethiconol Beeswax; Dimethiconol  
8 Behenate; Dimethiconol Borageate; Dimethiconol Dhupa Butterate; Dimethiconol  
9 Fluoroalcohol Dillnoleic Acid; Dimethiconol Hydroxystearate; Dimethiconol Illipe Butterate;  
10 Dimethiconol Isostearate; Dimethiconol Kokum Butterate; Dimethiconol Lactate; Dimethiconol  
11 Mohwa Butterate; Dimethiconol Sal Butterate; Dimethiconol Stearate; Dimethyl Adipate;  
12 Dimethylaminoethyl Methacrylate; Dimethyl Brassylate; Dimethyl Cystinate; Dimethyl  
13 Glutarate; Dimethyl Maleate; Dimethyl Oxalate; Dimethyl Phthalate; Dimethyl Succinate;  
14 Dimyristyl Tartrate; Dimyristyl Thiodipropionate; Dinonoxynol-9 Citrate; Dioctyl Adipate;  
15 Dioctyl Butamido Triazone; Dioctyl Dimer Dilinoleate; Dioctyldodeceth-2 Lauroyl Glutamate;  
16 Dioctyldodecyl Adipate; Dioctyldodecyl Dimer Dilinoleate; Dioctyldodecyl Dodecanedioate;  
17 Dioctyldodecyl Fluoroheptyl Citrate; Dioctyldodecyl Lauroyl Glutamate; Dioctyldodecyl  
18 Stearoyl Dimer Dilinoleate; Dioctyldodecyl Stearoyl Glutamate; Dioctyl Fumarate; Dioctyl  
19 Malate; Dioctyl Maleate; Dioctyl Phthalate; Dioctyl Sebacate; Dioctyl Succinate; Dioleoyl  
20 Edetolmonium Methosulfate; Dipalmitoyl Hydroxyproline; Dipentaerythrityl  
21 Hexacaprylate/Hexacaprate; Dipentaerythrityl Hexaheptanoate/Hexacaprylate/Hexacaprate;  
22 Dipentaerythrityl Hexahydroxystearate; Dipentaerythrityl

1 Hexahydroxystearate/Stearate/Rosinate; Dipentaerythrityl Hexaoctanoate/Behenate;  
2 Dipentaerythrityl Pentahydroxystearate/Isostearate; Diphenyl Carbomethoxy Acetoxy  
3 Naphthopyran; Dipropyl Adipate; Dipropylene Glycol Caprylate; Dipropylene Glycol  
4 Dibenzoate; Dipropylene Glycol Salicylate; Dipropyl Oxalate; Disodium Laureth-7 Citrate;  
5 Disodium PEG-5 Laurylcitrate Sulfosuccinate; Disodium PEG-8 Ricinosuccinate; Disodium  
6 Succinoyl Glycyrrhetinate; Disodium 2-Sulfolaurate; Disteareth-2 Lauroyl Glutamate;  
7 Disteareth-5 Lauroyl Glutamate; Distearyl Thiodipropionate; Ditallowoylethyl  
8 Hydroxyethylmonium Methosulfate; Ditridecyl Adipate; Ditridecyl Dimer Dilinoleate;  
9 Ditridecyl Thiodipropionate; Dodecyl Gallate; Erucyl Arachidate; Erucyl Erucate; Erucyl  
10 Oleate; Ethiodized Oil; Ethoxydiglycol Acetate; Ethoxyethanol Acetate; Ethyl Acetate; Ethyl  
11 Almondate; Ethyl Apricot Kernelate; Ethyl Arachidonate; Ethyl Aspartate; Ethyl Avocadate;  
12 Ethyl Benzoate; Ethyl Biotinate; Ethyl Butylacetylaminopropionate; Ethyl Cinnamate; Ethyl  
13 Cyanoacrylate; Ethyl Cyclohexyl Propionate; Ethyl Digydroxypropyl PABA; Ethyl  
14 Diisopropylcinnamate; Ethylene Brassylate; Ethylene Carbonate; Ethyl Ester of Hydrolyzed  
15 Animal Protein; Ethyl Ester of Hydrolyzed Keratin; Ethyl Ester of Hydrolyzed Silk; Ethyl  
16 Ester of PVM/MA Copolymer; Ethyl Ferulate; Ethyl Glutamate; Ethyl Isostearate; Ethyl  
17 Lactate; Ethyl Laurate; Ethyl Linoleate; Ethyl Linolenate; Ethyl Methacrylate; Ethyl  
18 Methoxycinnamate; Ethyl Methylphenylglycidate; Ethyl Minkate; Ethyl Morrhuate; Ethyl  
19 Myristate; Ethyl Nicotinate; Ethyl Oleate; Ethyl Olivinate; Ethyl PABA Ethyl Palmitate;  
20 Ethylparaben; Ethyl PCA; Ethyl Pelargonate; Ethyl Persate; Ethyl Phenylacetate; Ethyl  
21 Ricinoleate; Ethyl Serinate; Ethyl Stearate; Ethyl Thioglycolate; Ethyl Urocanate; Ethyl Wheat  
22 Germate; Ethyl Ximenynate; Etocrylene; Farnesyl Acetate; Galactonolactone; Galbanum

1 (Ferula Galbaniflua) Oil; Gamma-Nonalacione; Geranyl Acetate; Glucarolactone; Glucose  
2 Glutamate; Glucose Pentaacetate; Glucuronolactone; Glycereth-7 Benzoate; Glycereth-7  
3 Diisononanoate; Glycereth-8 Hydroxystearate; Glycereth-5 Lactate; Glycereth-25 PCA  
4 Isostearate; Glycereth-7 Triacetate; Glyceryl Triacetyl Hydroxystearate; Glyceryl Triacetyl  
5 Ricinoleate; Glycolamide Stearate; Glycol/Butylene Glycol Montanate; Glycol Catearate;  
6 Glycol Dibehenate; Glycol Dilaurate; Glycol Dioctanoate; Glycol Dioleate; Glycol Distearate;  
7 Glycol Ditallowate; Glycol Hydroxystearate; Glycol Montanate; Glycol Octanoate; Glycol  
8 Oleate; Glycol Palmitate; Glycol Ricinoleate; Glycol Salicylate; Glycol Stearate; Glycol  
9 Stearate SE; Glycyrrhetinyl Stearate; Hexacosyl Glycol Isostearate; Hexanediol Beeswax;  
10 Hexanediol Distearate; Hexanetriol Beeswax; Hexyldecyl Benzoate; Hexyldecyl Ester of  
11 Hydrolyzed Collagen; Hexyldecyl Isostearate; Hexyldecyl Laurate; Hexyldecyl Octanoate;  
12 Hexyldecyl Oleate; Hexyldecyl Palmitate; Hexyldecyl Stearate; Hexyldodecyl Salicylate;  
13 Hexyl Isostearate; Hexyl Laurate; Hexyl Nicotinate; Homosalate; Hydrogenated Castor Oil  
14 Hydroxystearate; Hydrogenated Castor Oil Isostearate; Hydrogenated Castor Oil Laurate;  
15 Hydrogenated Castor Oil Stearate; Hydrogenated Castor Oil Triisostearate; Hydrogenated  
16 Methyl Abietate; Hydrogenated Rosin; Hydroquinone PCA; Hydroxycetyl Isostearate;  
17 Hydroxyoctacosanyl Hydroxystearate; Inositol Hexa-PCA; Iodopropynyl Butylcarbamate;  
18 Isoamyl Acetate; Isoamyl Laurate; Isoamyl p-Methoxycinnamate; Isobutyl Acetate;  
19 Isobutylated Lanolin Oil; Isobutyl Benzoate; Isobutyl Myristate; Isobutyl Palmitate;  
20 Isobutylparaben; Isobutyl Pelargonate; Isobutyl Stearate; Isobutyl Tallowate; Isoceteareth-8  
21 Stearate; Isoceteth-10 Stearate; Isocetyl Behenate; Isocetyl Isodecanoate; Isocetyl Isostearate;  
22 Isocetyl Laurate; Isocetyl Linoleoyl Stearate; Isocetyl Myristate; Isocetyl Octanoate; Isocetyl

1 Palmitate; Isocetyl Salicylate; Isocetyl Stearate; Isocetyl Stearoyl Stearate; Isodeceth-2  
2 Cocoate; Isodecyl Citrate; Isodecyl Cocoate; Isodecyl Hydroxystearate; Isodecyl Isononanoate;  
3 Isodecyl Laurate; Isodecyl Myristate; Isodecyl Neopentanoate; Isodecyl Octanoate; Isodecyl  
4 Oleate; Isodecyl Palmitate; Isodecylparaben; Isodecyl Salicylate; Isodecyl Stearate; Isohexyl  
5 Laurate; Isohexyl Neopentanoate; Isohexyl Palmitate; Isolauryl Behenate; Isomerized Jojoba  
6 Oil; Isononyl Ferulate; Isooctyl Thioglycolate; Isopropyl Acetate; Isopropyl Arachidate;  
7 Isopropyl Avocadate; Isopropyl Behenate; Isopropyl Benzoate; Isopropylbenzyl Salicylate;  
8 Isopropyl Citrate; Isopropyl C12-15-Pareth-9 Carboxylate; Isopropyl Hydroxystearate;  
9 Isopropyl Isostearate; Isopropyl Jojobate; Isopropyl Lanolate; Isopropyl Laurate; Isopropyl  
10 Linoleate; Isopropyl Myristate; Isopropyl Oleate; Isopropylparaben; Isopropyl PPG-2-  
11 Isodeceth-7 Carboxylate; Isopropyl Ricinoleate; Isopropyl Sorbate; Isopropyl Stearate;  
12 Isopropyl Tallowate; Isopropyl Thioglycolate; Isosorbide Laurate; Isosteareth-10 Stearate;  
13 Isostearyl Avocadate; Isostearyl Behenate; Isostearyl Benzoate; Isostearyl Erucate; Isostearyl  
14 Isononanoate; Isostearyl Isostearate; Isostearyl Isostearoyl Stearate; Isostearyl Lactate;  
15 Isostearyl Laurate; Isostearyl Myristate; Isostearyl Neopentanoate; Isostearyl Octanoate;  
16 Isostearyl Palmitate; Isostearyl Stearoyl Stearate; Isotridecyl Isononanoate; Isotridecyl Laurate;  
17 Isotridecyl Myristate; Jojoba (Buxus Chinensis) Oil; Jojoba Esters; Kojic Dipalmitate; Laneth-  
18 9 Acetate; Laneth-10 Acetate; Laneth-4 Phosphate; Lanolin Linoleate; Lanolin Ricinoleate;  
19 Laureth-2 Acetate; Laureth-2 Benzoate; Laureth-6 Citrate; Laureth-7 Citrate; Laureth-2  
20 Octanoate; Laureth-7 Tartrate; Lauroyl Ethyl Glucoside; Lauroyl Lactic Acid; Lauryl  
21 Behenate; Lauryl Cocoate; Lauryl Isostearate; Lauryl Lactate; Lauryl Methacrylate; Lauryl  
22 Myristate; Lauryl Octanoate; Lauryl Oleate; Lauryl Palmitate; Lauryl Stearate; Linalyl

1 Acetate; Linoleyl Lactate; Madecassicoside; Mannitan Laurate; Mannitan Oleate; Menthyl  
2 Acetate; Menthyl Anthranilate; Menthyl Lactate; Menthyl PCA; Menthyl Salicylate;  
3 Methoxyisopropyl Acetate; Methoxy-PEG-7 Rutinyl Succinate; Methyl Acetate; Methyl Acetyl  
4 Ricinoleate; Methyl Anthranilate; Methyl Behenate; Methyl Benzoate; Methyl Caproate;  
5 Methyl Caprylate; Methyl Caprylate/Caprates; Methyl Cocoate; 6-Methyl Coumarin; Methyl  
6 Dehydroabietate; Methyl Dihydroabietate; Methyl dihydrojasmonate; Methyl Gluceth-20  
7 Benzoate; Methyl Glucose Dioleate; Methyl Glucose Isostearate; Methyl Glucose Laurate;  
8 Methyl Glucose Sesquicaprylate/Sesquicaprate; Methyl Glucose Sesquicocoate; Methyl  
9 Glucose Sesquiisostearate; Methyl Glucose Sesquilaurate; Methyl Glucose Sesquioleate;  
10 Methyl Glucose Sesquisteate; Methyl Glycyrrhizate; Methyl Hydrogenated Rosinate; Methyl  
11 Hydroxystearate; Methyl Isostearate; Methyl Laurate; Methyl Linoleate; Methyl 3-  
12 Methylresorcylate; Methyl Myristate; Methyl Nicotinate; Methyl Oleate; Methyl Palmate;  
13 Methyl Palmitate; Methylparaben; Methyl Pelargonate; Methyl Ricinoleate; Methyl Rosinate;  
14 Methyl Salicylate; Methylsilanol Acetylmethionate; Methylsilanol Carboxymethyl  
15 Theophylline; Methylsilanol Carboxymethyl Theophylline Alginate; Methylsilanol  
16 Hydroxyproline; Methylsilanol Hydroxyproline Aspartate; Methylsilanol Mannuronate;  
17 Methylsilanol PCA; Methyl Soyate; Methyl Stearate; Methyl Thioglycolate; Monosaccharide  
18 Lactate Condensate; Myreth-3 Caprate; Myreth-3 Laurate; Myreth-2 Myristate; Myreth-3  
19 Myristate; Myreth-3 Octanoate; Myreth-3 Palmitate; Myristoyl Ethyl Glucoside; Myristoyl  
20 Lactic Acid; Myristyl Isostearate; Myristyl Lactate; Myristyl Lignocerate; Myristyl  
21 Myristate; Myristyl Octanoate; Myristyl Propionate; Myristyl Salicylate; Myristyl Stearate;  
22 Neopentyl Glycol Dicaprate; Neopentyl Glycol Dicaprylate/Dicaprate; Neopentyl Glycol



1 Dicaprylate/Dipelargonate/Dicaprate; Neopentyl Glycol Diheptanoate; Neopentyl Glycol  
2 Diisostearate; Neopentyl Glycol Dilaurate; Neopentyl Glycol Dioctanoate; Nonyl Acetate;  
3 Nopyl Acetate; Octacosanyl Glycol Isostearate; Octocrylene; Octyl Acetoxystearate; Octyl  
4 Benzoate; Octyl Caprylate/Caprate; Octyl Cocoate; Octyldecyl Oleate; Octyldodecyl Behenate;  
5 Octyldodecyl Benzoate; Octyldodecyl Erucate; Octyldodecyl Hydroxystearate; Octyldodecyl  
6 Isostearate; Octyldodecyl Lactate; Octyldodecyl Lanolate; Octyldodecyl Meadowfoamate;  
7 Octyldodecyl Myristate; Octyldodecyl Neodecanoate; Octyldodecyl Neopentanoate;  
8 Octyldodecyl Octanoate; Octyldodecyl Octyldodecanoate; Octyldodecyl Oleate; Octyldodecyl  
9 Olivat; Octyldodecyl Ricinoleate; Octyldodecyl Stearate; Octyldodecyl Steroyl Stearate; Octyl  
10 Gallate; Octyl Hydroxystearate; Octyl Hydroxystearate Benzoate; Octyl Isononanoate; Octyl  
11 Isopalmitate; Octyl Isostearate; Octyl Laurate; Octyl Linoleayl Stearate; Octyl  
12 Methoxycinnamate; Octyl Myristate; Octyl Neopentanoate; Octyl Octanoate; Octyl Oleate;  
13 Octyl Palmitate; Octyl PCA; Octyl Pelargonate; Octyl Salicylate; Octyl Stearate; Oleoyl Ethyl  
14 Glucoside; Oleth-2 Benzoate; Oleyl Acetate; Oleyl Arachidate; Oleyl Erucate; Oleyl Ethyl  
15 Phosphate; Oleyl Lactate; Oleyl Lanolate; Oleyl Linoleate; Oleyl Myristate; Oleyl Oleate;  
16 Oleyl Phosphate; Oleyl Stearate; Oryzanol; Ozonized Jojoba Oil; Palmitoyl Carniline;  
17 Palmitoyl Inulin; Palmitoyl Myristyl Serinate; Pantethine; Panthenyl Ethyl Ester Acetate;  
18 Panthenyl Triacetate; PCA Glyceryl Oleate; Pea Palmitate; PEG-18 Castor Oil Dioleate; PEG-  
19 5 DEDM Hydantoin Oleate; PEG-15 DEDM Hydantoin Stearate; PEG-30  
20 Dipolyhydroxystearate; PEG-20 Hydrogenated Castor Oil Isostearate; PEG-50 Hydrogenated  
21 Castor Oil Isostearate; PEG-20 Hydrogenated Castor Oil Triisostearate; PEG-20 Mannitan  
22 Laurate; PEG-20 Methyl Glucose Distearate; PEG-80 Methyl Glucose Laurate; PEG-20


1 Methyl Glucose Sesquicaprylate/Sesquicaprate; PEG-20 Methyl Glucose Sesquilaurate; PEG-5  
2 Oleamide Dioleate; PEG-150 Pentaerythrityl Tetrastearate; PEG-3/PPG-2 Glyceryl/Sorbitol  
3 Hydroxystearate/Isostearate; PEG-4 Proline Linoleate; PEG-4 Proline Linolenate; PEG-8  
4 Propylene Glycol Cocoate; PEG-55 Propylene Glycol Oleate; PEG-25 Propylene Glycol  
5 Stearate; PEG-75 Propylene Glycol Stearate; PEG-120 Propylene Glycol Stearate; PEG-40  
6 Sorbitol Hexaoleate; PEG-50 Sorbitol Hexaoleate; PEG-30 Sorbitol Tetraoleate Laurate; PEG-  
7 60 Sorbitol Tetrastearate; PEG-5 Tricapryl Citrate; PEG-5 Tricetyl Citrate; PEG-5 Trilauryl  
8 Citrate; PEG-5 Trimethylolpropane Trimyristate; PEG-5 Trimyristyl Citrate; PEG-5 Tristearyl  
9 Citrate; PEG-6 Undecylenate; Pentadecalacione; Pentaerythrityl Dioleate; Pentaerythrityl  
10 Distearate; Pentaerythrityl Hydrogenated Rosinate; Pentaerythrityl  
11 Isostearate/Caprate/Caprylate/Adipate; Pentaerythrityl Rosinate; Pentaerythrityl Stearate;  
12 Pentaerythrityl Stearate/Caprate/Caprylate/Adipate; Pentaerythrityl Stearate/Isostearate/Adipate/  
13 Hydroxystearate; Pentaerythrityl Tetraabietate; Pentaerythrityl Tetraacetate; Pentaerythrityl  
14 Tetrabehenate; Pentaerythrityl Tetrabenzoate; Pentaerythrityl Tetracaprylate/Tetracaprate;  
15 Pentaerythrityl Tetracocoate; Pentaerythrityl Tetraisononanoate; Pentaerythrityl Tetralaurate;  
16 Pentaerythrityl Tetramyristate; Pentaerythrityl Tetraoctanoate; Pentaerythrityl Tetraoleate;  
17 Pentaerythrityl Tetrapelargonate; Pentaerythrityl Tetrastearate; Pentaerythrityl Trioleate;  
18 Phenethyl Acetate; Phenolphthalein; Phenoxyethylparaben; Phenyl Benzoate; Phenylparaben;  
19 Phenyl Salicylate; Phylosteryl Macadamiate; Poloxamer 105 Benzoate; Poloxamer 182  
20 Dibenzoate; Polycaprolactone; Polydimethylaminoethyl Methacrylate; Polyethylacrylate;  
21 Polyethylglutamate; Polyethylmethacrylate; Polymethyl Acrylate; Polymethylglutamate;  
22 Polysorbate 80 Acetate; Polyvinyl Acetate; Potassium Butylparaben; Potassium Deceth-4

1     Phosphate; Potassiu Ethylparaben; Potassium Methylparaben; Potassium Propylparaben; PPG-  
2     2 Isocelest-20 Acetate; PPG-14 Laureth-60 Alkyl Dicarbamate; PPG-20 Methyl Glucose Ether  
3     Acetate; PPG-20 Methyl Glucose Ether Distearate; PPG-2 Myristyl Ether Propionate; PPG-14  
4     Palmeth-60 Alkyl Dicarbamate; PPG-15 Steryl Ether Benzoate; Pregnenolone Acetate; Propyl  
5     Acetate; Propyl Benzoate; Propylene Carbonate; Propylene Glocol Alginate; Propylene Glycol  
6     Behenate; Propylene Glycol Caprylate; Propylene Glycol Ceteth-3 Acetate; Propylene Glycol  
7     Ceteth-3 Propionate; Propylene Glycol Citrate; Propylene Glycol Cocoate; Propylene Glycol  
8     Dicaprte; Propylene Glycol Dicaproate; Propylene Glycol Dicaprylate; Propylene Glycol  
9     Dicocoate; Propylene Glycol Diisononanoate; Propylene Glycol Diisostearate; Propylene  
10    Glycol Dilaurate; Propylene Glycol Dioctanoate; Propylene Glycol Dioleate; Propylene Glycol  
11    Dipelargonate; Propylene Glycol Distearate; Propylene Glycol Diundecanoate; Propylene  
12    Glycol Hydroxystearate; Propylene Glycol Isoceteth-3 Acetate; Propylene Glycol Isostearate;  
13    Propylene Glycol Laurate; Propylene Glycol Linoleate; Propylene Glycol Linolenate;  
14    Propylene Glycol Myristate; Propylene Glycol Myristyl Ether Acetate; Propylene Glycol  
15    Oleate; Propylene Glycol Oleate SE; Propylene Glycol Ricinoleate; Propylene Glycol Soyate;  
16    Propylene Glycol Stearate; Propylene Glycol Stearate SE; Propyl Gallate; Propylparaben;  
17    Pyricarbonate; Pyridoxine Dicaprylate; Pyridoxine Dilaurate; Pyridoxine Dioctenoate;  
18    Pyridoxine Dipalmitate; Pyridoxine Glycyrrhetinate; Pyridoxine Tripalmitate; Raffinose  
19    Myristate; Raffinose Oleate; Resorcinol Acetate; Retinyl Acetate; Retinyl Linoleate; Retinyl  
20    Palmitate; Retinyl Propionate; Riboflavin Tetraacetate; Ribonolactone; Rosin Acrylate;  
21    Siloxanetriol Phytate; Silybum Marianum Ethyl Ester; Sodium Behenoyl Lactylate; Sodium  
22    Butylparaben; Sodium Caproyl Lactylate; Sodium Cocoyl Lactylate; Sodium Dilaureth-7

1 Citrate; Sodium Ethylparaben; Sodium Ethyl 2-Sulfolaurate; Sodium Isostearoyl Lactylate;  
2 Sodium Laureth-7 Tartrate; Sodium Lauroyl Lactylate; Sodium Methylparaben; Sodium  
3 Methyl 2-Sulfolaurate; Sodium Oleoyl Lactylate; Sodium Panteheine Sulfonate; Sodium  
4 Phytate; Sodium Propylparaben; Sodium Stearoyl Lactylate; Sorbeth-2 Cocoate; Sorbeth-6  
5 Hexastearate; Sorbeth-3 Isostearate; Sorbityl Acetate; Soybean Palmitate; Soy Sterol Acetate;  
6 Stearamide DEA-Distearate; Stearamide DIBA-Stearate; Stearamide MEA-Stearate; Steareth-5  
7 Stearate; Stearoyl Lactylic Acid; Stearyl Acetate; Stearyl Acetyl Glutamate; Stearyl Beeswax;  
8 Stearyl Behenate; Stearyl Benzoate; Stearyl Caprylate; Stearyl Citrate; Stearyl Erucate; Stearyl  
9 Glycol Isostearate; Stearyl Glycyrrhetinate; Stearyl Heptanoate; Stearyl Lactate; Stearyl  
10 Linoleate; Stearyl Octanoate; Stearyl Stearate; Stearyl Stearoyl Stearate; Sucrose Acetate  
11 Isobutyrate; Sucrose Benzoate; Sucrose Cocoate; Sucrose Dilaurate; Sucrose Distearate;  
12 Sucrose Laurate; Sucrose Myristate; Sucrose Octaacetate; Sucrose Oleate; Sucrose Palmitate;  
13 Sucrose Polybehenate; Sucrose Polycottonseedate; Sucrose Polylaurate; Sucrose Polylinoleate;  
14 Sucrose Polypalmitate; Sucrose Polysoyate; Sucrose Polystearate; Sucrose Ricinoleate; Sucrose  
15 Stearate; Sucrose Tetrastearate Triacetate; Sucrose Tribehenate; Sucrose Tristearate; Tallowoyl  
16 Ethyl Glucoside; Tannic Acid; TEA-Lauroyl Lactylate; Telmestine; Terpeneol Acetate;  
17 Tetraethyl Phenyl Hydroxybenzoate; Tetradecyleicosyl Stearate; Tetrahexyldecyl Ascorbate;  
18 Tetrahydrofurfuryl Acetate; Tetrahydrofurfuryl Ricinoleate; Tocophersolan; Tocopheryl  
19 Acetate; Tocopheryl Linoleate; Tocopheryl Linoleate/Oleate; Tocopheryl Nicotinate;  
20 Tocopheryl Succinate; Tributyl Citrate; Tri-C12-13 Alkyl Citrate; Tri-C14-15 Alkyl Citrate;  
21 Tricaprylyl Citrate; Tridecyl Behenate; Tridecyl Cocoate; Tridecyl Erucate; Tridecyl  
22 Isononanoate; Tridecyl Laurate; Tridecyl Myristate; Tridecyl Neopentanoate; Tridecyl

1 Octanoate; Tridecyl Stearate; Tridecyl Stearoyl Stearate; Tridecyl Trimellitate; Triethyl  
2 Citrate; Triethylene Glycol hydrogenated Rosinate; Trihexyldecyl Citrate; Triisocetyl Citrate;  
3 Triisopropyl Citrate; Triisopropyl Trilinoleate; Triisostearyl Citrate; Triisostearyl Trilinoleate;  
4 Trilactin; Trilauryl Citrate; Trimethylolpropane Tricaprylate/Tricaprate; Trimethylolpropane  
5 Tricocoate; Trimethylolpropane Trilaurate; Trimethylolpropane Trioctanoate;  
6 Trimethylolpropane Tristearate; Trimethyl Pentanyl Diisobutyrate; Trioctyl Citrate;  
7 Trioctyldodecyl Borate; Trictyl Trimellitate; Trioleyl Citrate; TriPABA Panthenol;  
8 Tripropylene Glycol Citrate; Tristearyl Citrate; Tristearyl Phosphate; Vinyl Acetate; and Yeast  
9 Palmitate.

10 In addition to esters, alcohols may be gelled to form a gelled alcohol by using a diblock  
11 copolymer, a triblock copolymer, a star polymer, a radial polymer, a multi-block copolymer,  
12 and mixtures thereof. Suitable alcohols also may be gelled by alkyl galactomannan,  
13 polybutadiene, or other aforementioned polymers. Any alcohols as represented by the  
14 following formula may be gelled in embodiments of the invention.



17 where R represents any organic functional group which includes, but is not limited to,  
18 hydrocarbyl, phenyl, methoxyphenyl, and alkylphenyl, substituted alkyl, substituted phenyl,  
19 etc. Preferred alcohols include, but are not limited to, isostearyl alcohol, and octyl dodecanol.  
20 Other suitable alcohols include, but are not limited to, Abietyl Alcohol; Arachidyl Alcohol;  
21 Batyl Alcohol; Behenyl Alcohol; Benzyl Alcohol; Bishydroxyethyl Biscetyl Malonamide;  
22 Borneol; 2-t-Butylcyclohexyloxybutanol; Butyloctanol; C9-11 Alcohol; C12-13 Alcohol; C12-


1 15 Alcohol; C12-16 Alcohol; C14-15 Alcohol; C-20-40 Alcohols; C30-50 Alcohols; C40-60  
2 Alcohols; C18-38 Alkyl Hydroxystearoyl Stearate; Camphylcyclohexanol; Caproyl  
3 Sphingosine; Caprylic Alcohol; Caprylyl Glycol; CD Alcohol 19; Ceramide 1; Ceramide 2;  
4 Ceramide 3; Ceramide 4; Ceramide 5; Ceramide 1A; Ceramide 6 II; Cetearyl Alcohol; Cetyl  
5 Alcohol; Cetylarchidol; Cetyl Glycol; C9-13 Fluoroalcohol; C14-18 Glycol; C15-18 Glycol;  
6 C18-30 Glycol; C20-30 Glycol; Chimyl Alcohol; Chlorphenesin; Cholelciferol; Cholesterol;  
7 Cinnamyl Alcohol; Citronellol; Coconut Alcohol; Decyl Alcohol; Decyltetradecanol; 7-  
8 Dehydrocholesterol; Dichlorobenzyl Alcohol; Dihydrocholesterol; Dihydrolanosterol;  
9 Dihydroxyacetone; Dihydroxyethylamino Hydroxypropyl Oleate; 2,6-Dimethyl-7-Octen-2-ol;  
10 Dimethyl Octynediol; Dimethyl Phenylpropanol; Dodecylhexadecanol; Dodecyltetradecanol;  
11 Ergocalciferol; Ethyl Hexanediol; Farnesol; Galactonolactone; Geraniol; Glycyrrhetic Acid;  
12 Glycyrrhizic Acid; Heptylundecanol; Hexacosyl Glycol; 3-Hexenol; Hexyl Alcohol;  
13 Hexyldecanol; Hexyldecyloctadecanol; Hexylene Glycol; Hinokitiol; Hydroabietyl Alcohol;  
14 Hydrogenated Ethylbicycloneplane Guaiacol; Hydrogenated Tallow Alcohol; Hydrolyzed  
15 Glycyrrhizinate; Hydroxycapric Acid; Hydroxycaproyl Phytosphingosine; Hydroxycaprylic  
16 Acid; Hydroxycapryloyl Phytosphingosine; Hydroxyethyl Palmityl Oxyhydroxypropyl  
17 Palmitamide; Hydroxylauric Acid; Hydroxylauroyl Phytosphingosine; Hydroxymethyl  
18 Dioxoazabicyclooctane; Hydroxyproline; Hydroxystearyl Cetyl Ether; Jojoba Alcohol; Lactoyl  
19 Phytosphingosine; Lanolin Alcohol; Lauryl Alcohol; Lauryl Glycol; Linalool; p-Menthan-7-ol;  
20 Menthol; Menthone Glycerin Acetal; 3-Methylamino-4-Nitrophenoxyethanol; Methyl  
21 Glycyrrhizate; Methylsilanol Hydroxyproline; Myricyl Alcohol; Myristyl Alcohol; Nicotinyl  
22 Alcohol; Nicotinyl Tartrate; 3-Nitro-4-Aminophenoxyethanol; Octacosanyl Glycol;

1 Octoxyglycerin; Octoxyglyceryl Behenate; Octyldodecanol; 2-Oeamido-1,3-Octadecanediol;  
2 Oleyl Alcohol; Palm Alcohol; Palm Kernel Alcohol; Palmitamidohehexadecanediol; Panthenol;  
3 Panthenyl Ethyl Ether; Panthenyl Hydroxypropyl Steardimonium Chloride; Pentadecyl  
4 Alcohol; Pentylene Glycol; Phenethyl Alcohol; Phenoxyethanol; Phenoxyisopropanol;  
5 Phenylisohexanol; Phenylpropanol; Phytosphingosine; Polyvinyl Alcohol; Propanediol; Propyl  
6 Alcohol; Propylene Glycol; Pyridoxine Glycyrhetinate; Retinol; Ribonolactone; N-Stearoyl-  
7 Dihydroshingosine; Stearyl Alcohol; Stearyl Glycol; Tallow Alcohol; Terpeneol;  
8 Tetradecyleicosanol; Tetradecyloctadecanol; Tetrahydrofurfuryl Alcohol; Tetramethyl  
9 Cyclopentene Butenol; Tetramethyl Decynediol; Tridecyl Alcohol; Trimethylhexanol;  
10 Troxerutin; Undeceth-3; Undecylenyl Alcohol; and Undecylpentadecanol.

11 Numerous ethers also may be gelled to form gelled ether compositions by using a  
12 diblock copolymer, triblock copolymer, star polymer, radial polymer, multi-block copolymer,  
13 or mixtures thereof. Suitable ethers also may be gelled by alkyl galactomannan,  
14 polybutadiene, or other aforementioned polymers. Generally, an ether compound is  
15 represented by the following formula:



16  
17  
18  
19 where R and R' individually include, but are not limited to, hydrocarbyl, phenyl,  
20 methoxyphenyl, alkylphenyl, substituted alkyl, substituted phenyl, etc. Preferred ethers  
21 include, but are not limited to, dicarylyl ether and octyl methoxycinnamate. Dicarylyl ether  
22 is represented by the following chemical formula.




Other suitable ethers include, but are not limited to, Anethole; p-Anisic Acid; Batyl Alcohol; Batyl Isostearate; Batyl Stearate; Benzylhemiformal; 1,3-Bis-(2,4-Diaminophenoxy) propane; Butoxyethyl Acetate; Butoxyethyl Nicotinate; Butoxyethyl Stearate; Butoxypropanol; 2-t-Butylcyclohexyloxybutanol; Butyl Glucoside; Butylglucoside Caprate; Butyl Methoxydibenzoylmethane; Caprylyl/Capryl Glucoside; Capsaicin; Carboxymethyl Chitin; Carboxymethyl Chitosan Succinamide; Carboxymethyl Dextran; Cetearyl Glucoside; Cetyl Glyceryl Ether; Cetyl-PG Hydroxyethyl Decanamide; Cetyl-PG Hydroxyethyl Palmitamide; Chimyl Alcohol; Chimyl Isostearate; Chimyl Stearate; Chlorphenesin; Cinoxate; Cocamidopropyl Lauryl Ether; Coceth-4 Glucoside; Coco-Glucoside; Dibenzylidene Sorbitol; Dicetyl Ether; Dichlorophenyl Imidazoidioxolan; Dimethicone Copolyol Butyl Ether; Dimethicone Copolyol Ethyl Ether; Dimethicone Copolyol Methyl Ether; Dimethyl Hexahydronaphthyl Dihydroxymethyl Acetal; Dimethyl Isosorbide; Dioleyl Tocopheryl Methylsilanol; Diosmine; Disodium Cetyl Phenyl Ether Disulfonate; Disodium Decyl Phenyl Ether Disulfonate; Disodium Lauryl Phenyl Ether Disulfonate; Distarch Glyceryl Ether; Distearyl Ether; Ethoxydiglycol Acetate; Ethoxyheptyl Bicyclooctanone; 7-Ethylbicyclooxazolidine; Ethyl Methoxycinnamate; Ethyl Methylphenylglycidate; Ethyl Phenethyl Acetal; Eucalyptol; Eugenol; Ferulic Acid; Glyceryl Octanoate Dimethoxycinnamate; Glycofurol; Hexamethylindanopyran; Hexamidine; Hexamidine Diparaben; Hexamidine Paraben; Hydrogenated Ethylbicycloheptane Guaiacol; p-



1 Hydroxyanisole; Hydroxydecyl Maltitol; Hydroxyethyl Glyceryl Oleate/Stearate;  
2 Hydroxyethyl Palmityl Oxyhydroxypropyl Palmitamide; Hydroxyethyl Sorbitol;  
3 Hydroxymethoxybenzyl Pelargonamide; Hydroxypropyl Starch Phosphate; Hydroxystearyl  
4 Cetyl Ether; Isobutyl Methyl Tetrahydropyranol; Isoeugenol; Isolongifolene Epoxide;  
5 Isopropyl Hydroxycetyl Ether; Isostearamidopropyl Epoxypopyl Dimonium Chloride;  
6 Isostearyl Glyceryl Ether; Isostearyl Glyceryl Pentaerythrityl Ether; Lauryl Polyglyceryl-6  
7 Cetearyl Glycol Ether; Melatonin; Menthone Glycerin Acetal; Methoxyindane;  
8 Methoxyisopropyl Acetate; Methoxymethylbutanol; Methoxypropylgluconamide; Methylal;  
9 Ethyl Eugenol; Methyl Hexyl Ether; Methylsilanol Ascorbate; Myristyl-PG Hydroxyethyl  
10 Decanamide; 4-Nitroguaiacol; Octoxyglycerin; Octoxyglyceryl Behenate; Octoxyglyceryl  
11 Palmitate; Octyl Glyceryl Palmitate; Oleyl Glyceryl Ether; Panthenyl Ethyl Ether; Panthenyl  
12 Ethyl Ether Acetate; Panthenyl Hydroxypropyl Steardimonium Chloride; PEG-3 2,2-Di-p-  
13 Phenylenediamine; PEG-4 Ditallow Ether; PEG-150 Pentaerythrityl Tetrastearate; p-  
14 Phenetidine; Phenoxyethanol; Phenoxyethylparaben; Phenoxyisopropanol; Polyglycerin-3;  
15 Polyglycerin-4; Polyglycerin-6; Polyglycerin-10; Polyglyceryl-3 Cetyl Ether; Polyglyceryl-3  
16 Decyltetradecyl Ether; Polyglyceryl-3 Hydroxylauryl Ether; Polyglyceryl-2 Lanolin Alcohol  
17 Ether; Polyglyceryl-4 Lauryl Ether; Polyglyceryl-2 Oleyl Ether; Polyglyceryl-4 Oleyl Ether;  
18 Polyglyceryl Sorbitol; Polyvinyl Methyl Ether; Polyvinyl Stearyl Ether; PPG-9 Diglyceryl  
19 Ether; PPG-1-PEG-9 Lauryl Glycol Ether; Propylene Glycol Myristyl Ether; Quassin;  
20 Silanetriol Trehalose Ether; TEA-Lauryl Ether; Tetrahydrodiferuloyimethane; Thiodiglycol;  
21 Triclosan; Triethylene Glycol; Trihydroxypalmitamidohydroxypropyl Myristyl Ether;  
22 Trimethoxycaprylsilane; Troxerutin; and Ubiquinone.

Embodiments of the invention may also be used to obtain gelled natural products, such as naturally-occurring fats and oils. "Naturally-occurring fats and oils" often refer to the glyceryl esters of fatty acids (i.e., triglycerides) normally found in animal or plant tissues, including those which have been hydrogenated to reduce or eliminate unsaturation. Naturally occurring fats and oils include vegetable oils. Selected naturally-occurring fats and oils are represented by the following formula:



where R, R' and R'' may be the same or different fatty acid radicals.

Suitable naturally-occurring fats and oils that may be gelled by using a diblock copolymer, triblock copolymer, star polymer, radial polymer, multi-block copolymer or mixtures thereof. Suitable naturally-occurring fats and oils also may be gelled by alkyl galactomannan, polybutadiene, or other aforementioned polymers. Examples of the suitable naturally-occurring fats and oils include, but are not limited to, Adansonia Digitata Oil; Apricot (*Prunus Armeniaca*) Kernel Oil; *Argania Spinosa* Oil; *Argemone Mexicana* Oil;

- 1 Avocado (*Persea Gratissima*) Oil; Babassu (*Orbignya Oleifera*) Oil; Balm Mint (*Melissa*
- 2 *Officinalis*) Seed Oil; Bitter Almond (*Prunus Amygdalus Amara*) Oil; Bitter Cherry (*Prunus*
- 3 *Cerasus*) Oil; Black Currant (*Ribes Nigrum*) Oil; Borage (*Borago Officinalis*) Seed Oil; Brazil
- 4 (*Bertholletia Excelsa*) Nut Oil; Burdock (*Arctium Lappa*) Seed Oil; Butter; C12-18 Acid
- 5 Triglyceride; Calophyllum Tacamahaca Oil; Camellia Kissi Oil; Camellia Oleifera Seed Oil;
- 6 Canola Oil; Caprylic/Capric/Lauric Triglyceride; Caprylic/Capric/Linoleic Triglyceride;
- 7 Caprylic/Capric/Myristic/Stearic Triglyceride; Caprylic/Capric/Stearic Triglyceride;
- 8 Caprylic/Capric Triglyceride; Caraway (*Carum Carvi*) Seed Oil; Carrot (*Daucus Carota*
- 9 *Sativa*) Oil; Cashew (*Anacardium Occidentale*) Nut Oil; Castor Oil Benzoate; Castor (*Ricinus*
- 10 *Communis*) Oil; Cephalins; Chaulmoogra (*Taraktogenos Kurzii*) Oil, Chia (*Salvia Hispanica*)
- 11 Oil; Cocoa (*Theobroma Cacao*) Butter; Coconut (*Cocos Nucifera*) Oil; Cod Liver Oil; Coffee
- 12 (*Coffea Arabica*) Oil; Corn (*Zea Mays*) Germ Oil; Corn (*Zea Mays*) Oil; Cottonseed
- 13 (*Gossypium*) Oil; C10-18 Triglycerides; Cucumber (*Cucumis Sativus*) Oil; Dog Rose (*Rosa*
- 14 *Canina*) Hips Oil; Egg Oil; Emu Oil; Epoxidized Soybean Oil; Evening Primrose (*Oenothera*
- 15 *Biennis*) Oil; Fish Liver Oil; Gevuina Avellana Oil; Glyceryl Triacetyl Hydroxystearate;
- 16 Glyceryl Triacetyl Ricinoleate; Glycolipids; Glycosphingolipids; Goat Butter; Grape (*Vitis*
- 17 *Vinifera*) Seed Oil; Hazel (*Corylus Americana*) Nut Oil; Hazel (*Corylus Aveilana*) Nut Oil;
- 18 Human Placental Lipids; Hybrid Safflower (*Carthamus Tinctorius*) Oil; Hybrid Sunflower
- 19 (*Helianthus Annuus*) Seed Oil; Hydrogenated Canola Oil; Hydrogenated Castor Oil;
- 20 Hydrogenated Castor Oil Laurate; Hydrogenated Castor Oil Triisostearate; Hydrogenated
- 21 Coconut Oil; Hydrogenated Cottonseed Oil; Hydrogenated C12-18 Triglycerides;
- 22 Hydrogenated Fish Oil; Hydrogenated Lard; Hydrogenated Menhaden Oil; Hydrogenated Milk

1 Lipids; Hydrogenated Mink Oil; Hydrogenated Olive Oil; Hydrogenated Orange Roughy Oil;  
2 Hydrogenated Palm Kernel Oil; Hydrogenated Palm Oil; Hydrogenated Peanut Oil;  
3 Hydrogenated Rapeseed Oil; Hydrogenated Shark Liver Oil; Hydrogenated Soybean Oil;  
4 Hydrogenated Tallow; Hydrogenated Vegetable Oil; Isatis Tinctoria Oil; Job's Tears (Coix  
5 Lacryma-Jobi) Oil; Kiwi (Actinidia Chinensis) Seed Oil; Kukui (Aleurites Molluccana) Nut  
6 Oil; Lard; Lauric/Palmitic/Oleic Triglyceride; Linseed (Linum Usitatissimum) Oil; Lupin  
7 (Lupinus Albus) Oil; Macadamia Ternifolia Nut Oil; Maleated Soybean Oil; Mango  
8 (Mangifera Indica) Seed Oil; Marmot Oil; Meadowfoam (Limnanthes Alba) Seed Oil;  
9 Menhaden Oil; Milk Lipids; Mink Oil; Moringa Pterygosperma Oil; Mortierella Oil; Musk  
10 Rose (Rosa Moschata) Seed Oil; Neatsfoot Oil; Neem (Melia Azadirachta) Seed Oil; Oat  
11 (Avena Sativa) Kernel Oil; Oleic/Linoleic Triglyceride;  
12 Oleic/Palmitic/Lauric/Myristic/Linoleic Triglyceride; Oleostearine; Olive (Olea Europaea)  
13 Husk Oil; Olive (Olea Europaea) Oil; Omental Lipids; Orange Roughy Oil; Ostrich Oil;  
14 Oxidized Corn Oil; Palm (Elaeis Guineensis) Kernel Oil; Palm (Elaeis Guineensis) Oil;  
15 Passionflower (Passiflora Edulis) Oil; Peach (Prunus Persica) Kernel Oil; Peanut (Arachis  
16 Hypogaea) Oil; Pecan (Carya Illinoensis) Oil; Pengawar Djambi (Cibotium Barometz) Oil;  
17 Phospholipids; Pistachio (Pistacia Vera) Nut Oil; Placental Lipids; Poppy (Papaver Orientale)  
18 Oil; Pumpkin (Cucurbita Pepo) Seed Oil; Quinoa (Chenopodium Quinoa) Oil; Rapeseed  
19 (Brassica Campestris) Oil; Rice (Oryza Sativa) Bran Oil; Rice (Oryza Sativa) Germ Oil;  
20 Safflower (Carthamus Tinctorius) Oil; Salmon Oil; Sandalwood (Santalum Album) Seed Oil;  
21 Seabuchthorn (Hippophae Rhamnoides) Oil; Sesame (Sesamum Indicum) Oil; Shark Liver Oil;  
22 Shea Butter (Butyrospermum Parkii); Silk Worm Lipids; Skin Lipids; Soybean (Glycine Soja)

1 Oil; Soybean Lipid; Sphingolipids; Sunflower (*Helianthus Annuus*) Seed Oil; Sweet Almond  
2 (*Prunus Amygdalus Dulcis*) Oil; Sweet Cherry (*Prunus Avium*) Pit Oil; Tali Oil; Tallow; Tea  
3 Tree (*Melaleuca Alternifolia*) Oil; Telphairia Pedata Oil; Tomato (*Solanum Lycopersicum*)  
4 Oil; Triarachidin; Tribehenin; Tricaprin; Tricaprylin; Trichodesma Zeylanicum Oil; Trierucin;  
5 Triheptanoin; Triheptylundecanoin; Trihydroxymethoxystearin; Trihydroxystearin;  
6 Triisononanoin; Triisopalmitin; Triisostearin; Trilaurin; Trilinolein; Trilinolenin; Trimyristin;  
7 Trioctanoin; Triolein; Tripalmitin; Tripalmitolein; Triricinolein; Trisebacin; Tristearin;  
8 Triundecanoin; Tuna Oil; Vegetable Oil; Walnut (*Juglans Regia*) Oil; Wheat Bran Lipids; and  
9 Wheat (*Triticum Vulgare*) Germ Oil.

10 The gel compositions in accordance with embodiments of the invention may be  
11 obtained by the following method. First, one or more compounds to be gelled are mixed with  
12 a gelling agent. Second, the mixture typically is heated to a temperature in the range of about  
13 70° C. to about 140° C, although other temperatures also are acceptable. The mixture is  
14 agitated until a homogeneous mixture is obtained. The homogeneous mixture is then cooled  
15 to room temperature. A gel composition is thus obtained. It is noted that the compound to  
16 be gelled need not be mixed with a gelling agent before heating. Instead, the compound may  
17 be heated to a desired temperature first, and then the gelling agent is added to the compound.

18  
19 The following examples are given to illustrate embodiments of the invention. They are  
20 merely exemplary and are not intended to limit the scope of the invention otherwise described  
21 herein. All numerical values disclosed herein are approximate numbers. In some examples,  
22 an antioxidant was added in an amount of about 0.02%. Any antioxidants can be used. One

1 suitable antioxidant is 2,6-di-tert-butyl-4-methylphenol ("BHT").

2  
3 EXAMPLE 1

4 A sample of gelled isopropyl myristate (available under the trade name of Estol 1512,  
5 Lexol IPM, etc.) was prepared (85.28 wt. % isopropyl myristate + 14.30 wt. % Kraton G  
6 1702 + 0.40 wt. % Kraton G 1650). The finished gel exhibited excellent clarity and had a  
7 Brookfield viscosity (5 rpm, spindle T-C) of 157,000 CPS at 25° C.

8  
9 EXAMPLE 2

10 A sample of gelled octyl methoxycinnamate (trade names: Parsol MCX, Escalol 557,  
11 Neo Heliopan AV) was prepared (87.58 wt. % octyl methoxycinnamate + 12.00 wt. %  
12 Kraton G 1701 + 0.40 wt. % Kraton G 1650). The finished gel was clear and stable through  
13 several freeze/thaw cycles.

14  
15 EXAMPLE 3

16 A sample of zinc oxide suspension in gelled octyl methoxycinnamate was prepared by  
17 using 50.00 wt. % of zinc oxide and 50.00 wt. % of the gelled octyl methoxycinnamate  
18 illustrated in Example 2. The finished suspension had good consistency and maintained  
19 stability without separation for 15 days under thermal stress.

20  
21 EXAMPLE 4

22 A sample of gelled propylene glycol dicaprylate/caprate (trade name: Estol 1526) was

1 prepared (87.58 wt. % propylene glycol dicaprylate/caprate + 12.00 wt. % Kraton G 1780  
2 + 0.40 wt. % Kraton G-1650). The gel was clear at ambient temperature.

#### 3 4 EXAMPLE 5

5 A sample of gelled isostearyl neopentanoate (trade name: Dermol 185) was prepared  
6 (85.58 wt. % isostearyl neopentanoate + 14.00 wt. % Septon 1001 + 0.40 wt. % Kraton G  
7 1650). The finished gel was clear and stable without syneresis.

#### 8 9 EXAMPLE 6

10 Jojoba oil gel was prepared (91.28% jojoba oil + 0.4% Kraton G 1650 + 8.3% Kraton  
11 G 1702). The gel was clear and viscous.

#### 12 13 EXAMPLE 7

14 Eicosyl erucate (trade name: Erucical EG-20) gel was prepared (91.28% Erucical EG-  
15 20 + 0.4% Kraton G 1650 + 8.3% Kraton G 1702). The gel was clear and stable without  
16 syneresis.

17 In addition to the above examples of gel compositions, various other gel compositions  
18 also were obtained. Tables 3-6 summarize the gel compositions obtained in accordance with  
19 embodiments of the invention.

TABLE 3

INCI Name	Ester (wt.%)	Kraton® G1650	Kraton® G1702	Kraton® G1701	Kraton® G1780	Septon® 1001	Kraton® FG1901	Kraton® G1652
Isopropyl Myristate	85.28	0.4	14.3					
	90.58	0.4	9.00					
Isopropyl Palmitate	87.58	0.4	12.00					
	89.58	0.4	10.00					
	85.58	0.4	14.00					
C12-15 Alkyl Benzoate	83.58	0.4	16.00					
	87.58	0.4	12.00					
	87.98		12.00					
	89.58	0.4	10.00					
	89.98	10.00						
Octyl Methoxycinnamate	89.58	0.40	10.00					
	83.58	0.40	16.00					
	85.58	0.40	14.00					
	81.58	0.40	12.00					
	83.58	0.40		16.00				
	87.58	0.40		12.00				
	85.58	0.40		14.00				
Octyl Dodecyl Neopentanoate	89.58	0.40		10.00				
	81.98			12.00				
Isostearyl Neopentanoate	81.98	0.40				14.00		
	87.58	0.40	12.00					
Tridecyl Salicylate	91.28	0.4	8.30					
	89.58	0.4		10.00				
Octyl Dodecanol	88.89			6.67			4.44	
	90.00		5.48				4.5	
	90.00		5.45					4.5
	87.58	0.4	12.00					
	89.58	0.4	10.00					
Propylene Glycol Dicaprylate/caprate	87.58	0.4			12.00			
Jojoba Oil	91.28	0.4	8.30					



TABLE 4

	Polymer Type	Ester (%)	Polymer (%)	Antioxidant (%)
1,2-benzene-dicarboxylic acid, di-C <sub>8-10</sub> br alkyl ester	Vector® 6030	91.98	8	0.02
	Vector® 8550	91.98	8	0.02
	Vector® 2518P	91.98	8	0.02
	Solprene® S200	91.98	8	0.02
1,2 benzene-dicarboxylic acid, di-undecyl ester	Vector® 6030	91.98	8	0.02

TABLE 5

Oil Type	Polymer Type	Oil (%)	Polymer (%)	Antioxidant (%)
Sunflower Seed Oil	Kraton® D1102	91.9	8	0.10
	Kraton® D1133	91.9	8	0.10
	Kraton® D1101	91.9	8	0.10
	Vector® 6030	91.9	8	0.10
Corn Oil	Vector® 6030	91.9	8	0.10
Sesame Oil	Vector® 6030	91.9	8	0.10
Soybean Oil	Kraton® D1101	89.90	10	0.10
	Kraton® D1102	85.98	14	0.02
	Vector® 6030	91.90	8	0.10
Linseed Oil	Kraton® D1102	85.90	14	0.10
	Vector® 6030	91.90	8	0.10

TABLE 6

Oil Type	Oil (%)	Isopropyl Palmitate (%)	Alkyl Galactomannan (%)	Antioxidant (%)
Sunflower Seed Oil	46.99	46.99	6.0	0.02
	44.99	44.99	10.0	0.02
	75.18	18.8	6.0	0.02
	71.95	18.0	10.0	0.05
Corn Oil	75.1	18.8	6.0	0.10
	71.9	18.0	10.0	0.10
Soybean Oil	75.1	18.8	6.0	0.10
	71.9	18.0	10.0	0.10
Sesame Oil	68.9	25.0	6.0	0.10
	65.31	25.5	9.1	0.09
Olive Oil	68.9	25.0	6.0	0.10
	65.9	25.0	9.0	0.10

EXAMPLE 8

An *in-vivo* test was conducted to evaluate the waterproof efficacy of a sunscreen lotion based on a gelled ester composition. The *in-vivo* test procedure was conducted in accordance with the FDA monographs for sunscreen testing published in *Federal Register*, Vol. 43, No. 166, August 25, 1978. Briefly, a finished sunscreen lotion was applied to a skin surface which was subsequently submerged in water. Before water immersion, the sun protection factor ("SPF") was measured. After eighty minutes of water immersion, the SPF value was measured again. The percentage of the remaining SPF is indicative of the waterproof efficacy of the sunscreen lotion.

In this example, two sunscreen lotions were prepared according to Table 7. Both lotions contained approximately the same amount of a sunscreen active ingredient, i.e., octyl methoxycinnamate ("OMC"). Sunscreen Lotion A included gelled OMC, whereas Sunscreen Lotion B included neat OMC. Both sunscreen lotions contained approximately the same amounts of other ingredients as indicated in Table 7.

TABLE 7

Ingredient	Sunscreen Lotion A	Sunscreen Lotion B
De-ionized water	45.74	45.74
Tetra sodium EDTA	0.10	0.10
Propylene Glycol	2.50	2.50
Carbopol 940 (2% solution)	25.00	25.00
DEA-cetyl phosphate	1.00	1.00
Cetearyl Alcohol and Ceteareth 20	1.50	1.50
Mineral oil	—	1.26
Octyl Methoxycinnamate	—	7.50
Octyl Methoxycinnamate + Hydrogenated Ethylene/propylene/styrene copolymers + Hydrogenated Butylene/ethylene/styrene copolymers	8.76	—
Dimethicone	2.00	2.00
Diisodecyl Adipate	7.00	7.00
C <sub>12-15</sub> Alkyl Benzoate	5.00	5.00
Triethanol amine	0.40	0.40
Diazolidinyl urea + propylene glycol + methylparaben + propylparaben	1.00	1.00
Remaining SPF after 80 min. water immersion	84%	69%

After eighty minutes of water immersion, the remaining SPF for Sunscreen Lotion A was about 84% of the pre-immersion SPF. As to Sunscreen Lotion B, the remaining SPF was about 69% of its pre-immersion SPF. Therefore, the test result indicates that inclusion of a gelled sunscreen active ingredient increases the waterproof efficacy of the sunscreen lotion.

**EXAMPLE 9**

This example shows that gelled esters are comparable to petrolatum as an occlusive agent. An *in-vivo* skin barrier test was utilized to evaluate a gelled ester's ability to enhance the barrier function of the stratum corneum of human skin. Three different ester gels were formulated and evaluated in this test. Their respective formulations are listed in Table 8.

**TABLE 8**

Sample	Intermediate	Ester (wt. %)	Kraton G1701 (wt. %)	Kraton G1650 (wt. %)	Kraton G1701 (wt. %)	Anti- oxidant (%)
Gel No. 1	C <sub>12-15</sub> Alkyl Benzoate	85.58	14.00	0.40	—	0.02
Gel No. 2	Isopropyl Palmitate	87.58	12.00	0.40	—	0.02
Gel No. 3	Octyl Methoxycinnamate	85.58	—	0.40	14.00	0.02

An independent laboratory conducted Trans-Epidermal Water Loss ("TEWL") testing using the above gel compositions. Thirteen subjects were employed. TEWL measurements were taken on each subject at baseline and at 1, 2, and 3 hours after the application of a test material. A Servomed® Evaporimeter was used to measure the effectiveness of a topically applied test material. The evaporimeter samples relative humidity at two points above the skin surface which allowed the rate of water loss to be calculated from the measured humidity gradient. Petrolatum also was included in the measurements. The data obtained from the TEWL measurements are presented in Table 9.

TABLE 9

Test Material	Baseline	Hour 1		Hour 2		Hour 3	
		mean	$\Delta\%$	mean	$\Delta\%$	mean	$\Delta\%$
Gel No. 1	7.07	5.47	22.6%	5.44	23.1%	5.26	25.6%
Gel No. 2	6.66	5.37	19.4%	5.14	22.8%	4.72	29.1%
Gel No. 3	6.76	5.79	14.3%	5.49	18.8%	5.25	22.3%
Petrolatum	6.64	4.64	30.1%	4.28	35.5%	4.13	37.8%

Note:  $\Delta\% = (\text{Baseline} - \text{Mean}) * 100 / \text{Baseline}$ .

The data indicate that all three gelled esters are effective in retaining moisture in the skin. Consequently, gelled esters can be used as an occlusive agent in various health and beauty aid compositions.

#### EXAMPLE 10

This example shows that a gelled composition has significantly higher loading capacity as a suspension system. To compare the loading capacity of gelled soybean oil with that of neat soybean oil, two experiments were conducted: about 3 grams coated  $\text{TiO}_2$  powder were suspended in about 97 grams of neat soybean oil; and about 50 grams of the same coated  $\text{TiO}_2$  powder were suspended in 50 grams of gelled soybean oil. No dispersant agent was used in either suspension system. The gelled soybean oil was made from the following formulation: about 85.9 wt.% of commercially available soybean oil and about 14 wt.% of Kraton® D1102. It was observed that only about 2 wt.% of  $\text{TiO}_2$  remained suspended in the neat soybean oil. In contrast, about 50 wt.% of  $\text{TiO}_2$  was suspended in the gelled soybean oil. It is surprising that the loading capacity of gelled soybean oil is about 25 times more than that of neat soybean oil. The increased loading capacity by using a gelled naturally-occurring fat or oil should result in improved performance and cost savings.

1           Accordingly, the gel compositions in accordance with embodiments of the invention have  
2           numerous cosmetic and industrial applications. The gel compositions may be used alone or in  
3           combination with one or more additional ingredients. For example, the gel compositions are  
4           suitable as lubricants, suspending agents, emulsion stabilizers, thickening agents, personal care  
5           ingredients, air freshener components, pesticide and insecticide components, ingredients for candle  
6           and ornamental products, pharmaceutical carriers and carrier ingredients, ointment base  
7           ingredients, sporting goods, and vehicles for carrying other materials. Since esters are generally  
8           considered as biodegradable materials, this allows gelled esters to find application in products  
9           where minimal pollution is desired. Examples include fishing line lubricant, solder flux,  
10          agricultural dust reduction and lubrication, textile coating, protective coating for transporting  
11          fragile or environmentally-sensitive materials, and biodegradable oils and greases.

12          It has been discovered that the gel compositions in accordance with embodiments of the  
13          invention keep solids and/or liquids substantially uniformly suspended and evenly dispersed over  
14          a substantial period of time. The suspended liquids and/or solids may be present in the gel in  
15          amounts of up to about 95 weight percent. Suitable solids and/or liquids that can be suspended  
16          in the gel include any solid or non-hydrocarbon oil liquid which will disperse into the gel and  
17          remain substantially suspended or evenly dispersed therein.

18          Examples of suitable solids include, but are not limited to, zinc oxide, coated zinc oxide,  
19          surface-treated zinc oxide, titanium dioxide, coated titanium dioxide, surface-treated titanium  
20          dioxide, phosphorescent substances, fluorescent materials, molybdenum oxide, zinc sulfide,  
21          copper-doped zinc sulfide, graphite, explosive materials, pesticides, herbicides, fungicides,  
22          insecticides, plasticizers, air sensitive chemicals, moisture sensitive chemicals, boron nitride, iron

1 oxides, talc, mica, plastics, polymers, silica, silicon dioxide, aluminum oxide, inorganic materials,  
2 organometallic materials, metal particles, medical materials (such as antibacterials, antibiotics,  
3 antimicrobials, antifungals, and anesthetics), glass, clays, gums, capsules containing an active  
4 ingredient, starch, modified starch, other encapsulated materials, and mixtures thereof.

5 Examples of oil insoluble liquids which can be suspended in the gels include, but are not  
6 limited to, water, water containing one or more water soluble materials, glycerin, propylene  
7 glycol, butylene glycol, alcohols, acids, surfactants, emulsifiers, polyglycerols, ethers, polar esters,  
8 fluorinated compounds, perfluoropolyethers, silicones, silicon-containing compounds, and the  
9 mixtures thereof.

10 Pending U.S. Patent Application Serial No. 09/007,838, entitled "Hydrocarbon Gels as  
11 Suspending and the Dispersing Agents and Products," filed January 15, 1998, discloses methods  
12 of making a suspension system based on gelled hydrocarbons. The disclosed methods can be  
13 utilized in embodiments of the invention to make suspension systems which are based on gelled  
14 esters, gelled alcohols, gelled naturally-occurring fats and oils, or gelled ethers. The disclosure  
15 of the above-referenced patent application is incorporated by reference in its entirety herein.

16 The gel compositions in accordance with embodiments of the invention also have a wide  
17 spectrum of cosmetic applications when the gel compositions include an effective amount of one  
18 or more cosmetic and health and beauty aid ingredients. By "effective amount," it is meant that  
19 a sufficient amount of the ingredient is present to be effective for the indicated purpose in the  
20 composition. An effective amount may range from about 0.001 to about 80 wt.%. By "cosmetic  
21 and health and beauty aid ingredients," it is meant any material which can be applied topically to  
22 the human skin or any part thereof for cleansing, beautifying, promoting attractiveness, and

1 protecting or altering the appearance of the skin without altering or interfering with the  
2 physiological competence of the human skin or body. Included within this definition are creams,  
3 lotions, emollients, fragrance oils, moisturizers, humectants, cosmetic oils, and so on. The gel  
4 compositions in accordance with embodiments of the invention also may contain skin care  
5 preservatives, diluents, surfactants, anti-wrinkle agents, and the like.

6 Furthermore, the gel compositions may be utilized to manufacture various over-the-  
7 counter ("OTC") products. An OTC product may be made entirely from a gel composition, or  
8 only a component of the OTC is made from the gel composition. Examples of OTC products  
9 include, but are not limited to, antiperspirants, lip balms, and sunscreen (e.g., natural sunblocks,  
10 such as submicron particles of metal oxides or synthetic sunblock agents, such as octyl  
11 methoxycinnamate and benzophenone-3).

12 As mentioned above, the gel composition in accordance with embodiments of the  
13 invention may be utilized as carrier vehicles for topical administration of various cosmetic and  
14 health and beauty aid materials to the skin. Thus, such materials can be incorporated into the gel  
15 which are applied to the skin to be absorbed, to form a film on the skin, to provide a cooling  
16 sensation, to treat dry skin or oily skin, to work a material into the skin, to alter the overall texture  
17 of the skin, or to change color. All of these effects are sought to be achieved by various health  
18 and beauty aid products. Methods for making such skin care products are known in the art. For  
19 example, U.S. Patent No. 5,558,872 discloses a clear gelled mineral oil based skin protectant.  
20 Similar skin protectants may be made by substituting the mineral oil gel by the gel compositions  
21 in accordance with embodiments of the invention. The disclosure of U.S. Patent No. 5,558,872  
22 is incorporated by reference in its entirety herein.



1        It is noted that preferred uses of the gel compositions in accordance with embodiments  
2 of the invention include formation of thickened liquids, soft gels and semi-solid gels. Gels are  
3 particularly useful in waterproofing sunscreen compositions, makeup, mascara, etc. They also are  
4 useful in petrolatum-based products, such as petroleum jelly, makeup foundation, and night  
5 creams. They also can be used as substitutes for water-soluble polymers in products, such as lip  
6 rouge-cream, eyeliner liquid, and the like. They also may be used as a gelling agent in facial oil.

7        Semi-solid or solid gels have applications as toiletry sticks, such as a stick insect repellent  
8 or a matrix for clear or opaque stick products, which include deodorants, antiperspirants, lipsticks,  
9 analgesics, blushers, solid lotions and solid absorbable flexible gels. Methods for making such  
10 cosmetic sticks are known. For example, U.S. Patent No. 5,756,082 discloses a cosmetic stick  
11 composition based on a hydrocarbon oil gel. The disclosed hydrocarbon oil gels may be  
12 substituted by the gel compositions in accordance with embodiments of the invention to make  
13 cosmetic stick compositions. The disclosure of U.S. Patent No. 5,756,082 is incorporated by  
14 reference in its entirety herein.

15        As demonstrated above, the gel compositions in accordance with embodiments of the  
16 invention have a wide range of industrial and cosmetic applications. When so used, the gel  
17 compositions may exhibit one or more of the following properties or advantages: transparency;  
18 compatibility with active ingredients; reduction or elimination of syneresis; ability to act as a  
19 vehicle to provide a stable suspension of an ingredient; moisturization; enhancement of wash-off  
20 resistance; provision of improved SPF when formulated into sunscreen products; reduction in  
21 absorption and irritation; elimination or minimization of whitening effect; ability to act as a  
22 cosmetic base; controlled release of volatile ingredients; and formulation with less emulsifiers.

1 Other properties and advantages are apparent to a person of ordinary skill in the art.

2 While the invention has been described with respect to a limited number of embodiments,  
3 modifications and variations therefrom exist. For example, although suitable esters, ethers,  
4 alcohols, and naturally-occurring fats and oils have been described with some depth, other  
5 compounds also can be used. Additional suitable esters may include alkoxylated fatty acids,  
6 glyceryl ethers, and sorbitan derivatives. Additional suitable alcohols may include alkanolamides,  
7 alkanolamines, fatty alcohols, polyols, phenols, and sterols. Additional suitable ethers include  
8 heterocyclic ethers, e.g., tocopherol, alkoxylated alcohols, alkoxylated amides, alkoxylated  
9 amines, alkoxylated carboxylic acids, polymeric ethers, and glyceryl ethers. Similarly, naturally-  
10 occurring fats and oils may include so-called "essential oils," which are volatile organic  
11 constituents of plants normally obtained by distillation. In addition to the above-described  
12 compounds, aldehydes and ketones also may be formed into gel compositions in accordance with  
13 embodiments of the invention. The appended claims are intended to encompass all such  
14 modifications and the variations as falling within the scope of the invention.


15

16 What is claimed is:


1. A gel composition, comprising:  
 an ester compound; and  
 a polymer compound selected from the group consisting of triblock copolymers, star polymers, radial polymers, multi-block copolymers, and combinations thereof.

2. The gel composition of claim 1, further comprising a diblock copolymer,  
 wherein the gel composition is substantially free of mineral oils.

3. The gel composition of claim 1, wherein the ester is represented by the following formulas:




and



wherein  $n=1, 2, 3$ , and  $4$ , and  $\text{R}_1$  includes hydrogen, hydrocarbyl, phenyl, methoxyphenyl, alkylphenyl, substituted alkyl, and substituted phenyl;


$\text{R}_2$  includes hydrogen, hydrocarbyl, phenyl, methoxyphenyl, alkylphenyl, substituted alkyl, substituted phenyl, alkylene, phenylene, substituted alkylene, and substituted phenylene.

4. The gel composition of claim 1, wherein the ester is represented by the following formula:



wherein  $\text{R}_1$  includes hydrogen, hydrocarbyl, phenyl, methoxyphenyl, alkylphenyl, substituted alkyl, and substituted phenyl, and  $\text{R}_3$  includes alkylene, phenylene, substituted alkylene, or substituted phenylene.

5. The gel composition of claim 1, wherein the ester is represented by the following formula:



wherein  $\text{R}_4$ ,  $\text{R}_5$ , and  $\text{R}_6$  individually include alkylene, phenylene, substituted alkylene, or substituted phenylene, and  $\text{R}_7$ ,  $\text{R}_8$  and  $\text{R}_9$  individually include hydrogen, hydrocarbyl, phenyl, methoxyphenyl, alkylphenyl, substituted alkyl, and substituted phenyl.

6. The gel composition of claim 1, wherein the ester is selected from the group consisting of isopropyl myristate, isopropyl palmitate,  $\text{C}_{12}$ - $\text{C}_{15}$  alkyl benzoate, octyl methoxycinnamate, octyl dodecyl neopentanoate, propylene glycol dicaprylate/caprate, jojoba oil, and isostearyl neopentanoate.

7. The gel composition of claim 1, wherein the polymer compound is present in the amount of about 1% to about 40 % by weight.

8. The gel composition of claim 2, wherein the diblock copolymer is selected from the group consisting of styrene-ethylene/propylene copolymers, styrene-ethylene/butadiene copolymers, styrene-isoprene copolymers, and styrene-butadiene copolymers.

9. The gel composition of claim 1, wherein the triblock copolymer is selected from the group consisting of styrene-ethylene/propylene-styrene copolymers, styrene-ethylene/butadiene-styrene copolymers, styrene-isoprene-styrene copolymers, and styrene-butadiene-styrene copolymers.

- 1     10.     The gel composition of claim 2, wherein the diblock copolymer is hydrogenated.
- 1     11.     The gel composition of claim 1, wherein the triblock copolymer is hydrogenated.
- 1     12.     The gel composition of claim 1, wherein the triblock copolymer includes a grafted  
2 functional group.
- 1     13.     The gel composition of claim 1, further comprising a suspended component.
- 1     14.     The gel composition of claim 13, further comprising a diblock copolymer.
- 1     15.     The gel composition of claim 13, wherein the suspended component is a solid selected  
2 from the group consisting of organic materials, inorganic materials, organometallic materials,  
3 phosphorescent materials, and fluorescent materials.
- 1     16.     The gel composition of claim 13, wherein the suspended component is a solid selected  
2 from the group consisting of zinc oxide, coated zinc oxide, surface-treated zinc oxide, titanium  
3 dioxide, surface-treated titanium dioxide, graphite, explosives, air-sensitive chemicals, moisture-  
4 sensitive chemicals, boron nitride, iron oxides, talc, mica, plastics, polymers, silica, silicon dioxide,  
5 aluminum oxide, metal particles, antibacterials, antibiotics, anesthetics, glass, clays, gums,  
6 capsules containing an active ingredient, starch, modified starch, other encapsulated materials, and  
7 combinations thereof.
- 1     17.     The gel composition of claim 13, wherein the suspended component is a liquid selected  
2 from the group consisting of water, water containing a water-soluble material, glycerin, propylene  
3 glycol, butylene glycol, alcohols, acids, surfactants, emulsifiers, polyglycerols, ethers, polar esters,  
4 fluorinated compounds, perfluoropolyethers, silicones, silicon-containing compounds, and  
5 combinations thereof.
- 1     18.     The gel composition of claim 1, further comprising an active ingredient.

1 19. The gel composition of claim 18, wherein the active ingredient is selected from the group  
2 consisting of sunscreens, antiperspirants, deodorants, perfumes, cosmetics, emollients, insect  
3 repellants, pesticides, herbicides, fungicides, plasticizers, insecticides, and medicaments.

1 20. A gel composition, comprising:  
2 a compound selected from the group consisting of alcohols, ethers, naturally occurring fats  
3 and oils, and combinations thereof; and  
4 a polymer compound selected from the group consisting of diblock copolymers, triblock  
5 copolymers, star polymers, radial polymers, multi-block copolymers, and combinations thereof.

1 21. The gel composition of claim 20, wherein the alcohols include octyl dodecanol or  
2 isostearyl alcohol.

1 22. The gel composition of claim 20, wherein the ethers include dicarylyl ether.

1 23. The gel composition of claim 20, wherein the naturally occurring fats and oils include  
2 linseed oil, soybean oil, sunflower seed oil, corn oil, sesame oil, olive oil, castor oil, coconut oil,  
3 palm oil, and peanut oil.

1 24. A gel composition, comprising:  
2 a compound selected from the group consisting of esters, alcohols, ethers, naturally  
3 occurring fats and oils, and combinations thereof; and  
4 a polymer compound selected from the group consisting of alkyl galactomannan,  
5 polybutadiene, and combinations thereof.

- 1     25.     A method of making a gel composition, comprising:  
2             mixing an ester compound with a polymer compound selected from the group consisting  
3     of triblock copolymers, star polymers, radial polymers, multi-block copolymers, and combinations  
4     thereof,  
5             heating the mixture;  
6             agitating the mixture until the mixture becomes homogeneous; and  
7             cooling the mixture.
- 1     26.     A method of making a gel composition, comprising:  
2             mixing an alcohol, an ether or a naturally occurring fat or oil with a polymer compound  
3     selected from the group consisting of diblock copolymers, triblock copolymers, star polymers,  
4     radial polymers, multi-block copolymers, and combinations thereof,  
5             heating the mixture;  
6             agitating the mixture until the mixture becomes homogeneous; and  
7             cooling the mixture.
- 1     27.     A method of making a gel composition, comprising:  
2             mixing an ester, an alcohol, an ether or a naturally occurring fat or oil with alkyl  
3     galactomannan or polybutadiene,  
4             heating the mixture;  
5             agitating the mixture until the mixture becomes homogeneous; and  
6             cooling the mixture.

1/3



Diblock Copolymer

FIG. 1




Triblock Copolymer

FIG. 2




2/3



Radial Copolymer

FIG. 3



Star Copolymer

FIG. 4

3/3



Multi-block Copolymer

FIG 5A



Multi-block Copolymer

FIG. 5B

# INTERNATIONAL SEARCH REPORT

Intern. Application No  
PCT/US 99/24840

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C08K5/103

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C08K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 91 05014 A (RAYCHEM LTD) 18 April 1991 (1991-04-18)  claim 1; examples A,B	1,3, 7-11,19, 20,23,24
X	US 5 412 022 A (ANDRES JOHANNES ET AL) 2 May 1995 (1995-05-02)  column 3, line 25-61; examples 1,2 column 4, line 15-20	1,3, 7-15,20, 23,24
X	EP 0 210 655 A (NIPPON STEEL CHEMICAL CO) 4 February 1987 (1987-02-04) claims 7,8	1,3,5,7, 19
A	EP 0 147 146 A (AMERICAN HOME PROD) 3 July 1985 (1985-07-03) claims; examples	1-10

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

15 February 2000

Date of mailing of the international search report

24/02/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5518 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Friederich, P

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/24840

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9105014 A	18-04-1991	AT 133987 T	15-02-1996
		AU 638287 B	24-06-1993
		AU 6508190 A	28-04-1991
		CA 2065730 A	06-04-1991
		DE 69025331 D	21-03-1996
		DE 69025331 T	10-10-1996
		EP 0494936 A	22-07-1992
		JP 2971133 B	02-11-1999
		JP 5500976 T	25-02-1993
		US 5442004 A	15-08-1995
US 5412022 A	02-05-1995	DE 4039899 A	17-06-1992
		AT 138086 T	15-06-1996
		DE 59107825 D	20-06-1996
		DK 561843 T	14-10-1996
		WO 9210537 A	25-06-1992
		EP 0561843 A	29-09-1993
		ES 2086555 T	01-07-1996
		JP 6503105 T	07-04-1994
EP 0210655 A	04-02-1987	JP 1444042 C	08-06-1988
		JP 62030140 A	09-02-1987
		JP 62054819 B	17-11-1987
		DE 3682454 A	19-12-1991
		US 4742102 A	03-05-1988
EP 0147146 A	03-07-1985	AU 3683484 A	27-06-1985
		CA 1238275 A	21-06-1988
		DK 626384 A	23-06-1985
		GR 81250 A	19-11-1985
		HU 37040 A	28-11-1985
		JP 60152413 A	10-08-1985
		KR 8900183 B	09-03-1989
		US 4931283 A	05-06-1990
		US 4933184 A	12-06-1990
		ZA 8409780 A	27-08-1986